Cattaneo-type subdiffusion equation

Tadeusz Kosztołowicz*

Institute of Physics, Jan Kochanowski University, University 7, 25-406 Kielce, Poland and Department of Radiological Informatics and Statistics,

Medical University of Gdańsk, Tuwima 15, 80-210 Gdańsk, Poland

Aldona Dutkiewicz[†]

Faculty of Mathematics and Computer Science, Adam Mickiewicz University, Universytetu Poznańskiego 4, 61-614 Poznań, Poland

Katarzyna D. Lewandowska[‡]

Department of Physics and Biophysics, Medical University of Gdańsk, Dębinki 1, 80-211 Gdańsk, Poland (Dated: September 20, 2024)

The ordinary subdiffusion equation, with fractional time derivatives of at most first order, describes a process in which the propagation velocity of diffusing molecules is unlimited. To avoid this non-physical property the Cattaneo diffusion equation has been proposed. Compared to the ordinary subdiffusion equation, the Cattaneo equation contains an additional time derivative of order greater than one and less than or equal to two. The fractional order of the additional derivative depends on the subdiffusion exponent. We study a Cattaneo-type subdiffusion equation (CTSE) that differs from the ordinary subdiffusion equation by an additional integro-differential operator (AO) which may be independent of subdiffusion parameters. The AO describes processes affecting ordinary subdiffusion. The equation is derived combining the modified diffusive flux equation with the continuity equation. It can also be obtained within the continuous time random walk model with the waiting time distribution for the molecule to jump controlled by the kernel of AO. We discuss whether the ordinary subdiffusion equation and CTSE provide qualitative differences in the description of subdiffusion. For example, we consider two types of CTSE: the CTSE with AO which is the Caputo fractional time derivative of the order independent of the subdiffusion exponent and with the AO with a kernel that is a slowly varying function. In the first case the effect generated by AO disappears relatively quickly over time. In the second one the effect may be visible for a long time.

PACS numbers:

I. INTRODUCTION

Subdiffusion occurs in media in which particle random walk is very hindered. The examples are transport of molecules in viscoelastic chromatin network [1], porous media [2], living cells [3], transport of sugars in agarose gel [4], and antibiotics in bacterial biofilm [5]. Subdiffusion is often defined as an anomalous diffusion process in which the temporal evolution of mean square displacement (MSD) of a particle is $\sigma^2(t) = 2Dt^{\alpha}/\Gamma(1+\alpha)$, where $\alpha \in (0, 1)$ is the subdiffusion parameter (exponent), D is a subdiffusion coefficient given in units of m^2/s^{α} . Frequently, ordinary subdiffusion is described by a fractional differential equation [6–14] with the Riemann-Liouville fractional time derivative of $1 - \alpha$ order or an equation with the Caputo fractional time derivative of the order α .

For the parabolic normal diffusion equation and subdiffusion equation with a time derivative of at most first order the Green's function P(x, t), which is the probability density of finding the molecule at position x at time t, is greater than zero for any x and t > 0. It means that the propagation velocity of some particles is arbitrarily high. To avoid this non-physical property, the Cattaneo diffusion equation has been proposed; the equation was originally used to describe heat propagation [15]. The equation involves a second-order time derivative controlled by the parameter τ , see Eq. (16) presented later. One of the models that provide the equation is based on the assumption that the flux of diffusing molecules is delayed by time τ with respect to the concentration gradient.

While the Cattaneo hyperbolic diffusion equation is well defined, the Cattaneo subdiffusion equation can take various non-equivalent forms, see for example Refs. [16– 24]. It contains a time derivative of the order greater than one and less than or equal to two; the fractional order of this derivative depends on α . The Cattaneo equation with the Caputo and/or the Riemann–Liouville fractional derivatives have been used to describe subdiffusion and subdiffusion with reactions [25–30], in particular neutron transport inside the core of a nuclear reactor [31], heat transport in porous media [32], and in a system with glass spheres in a tank filled with air [33]. Cattaneo equations with fractional derivatives other than those mentioned

^{*}Electronic address: tadeusz.kosztolowicz@ujk.edu.pl

 $^{^{\}dagger} \rm Electronic ~address: szukala@amu.edu.pl$

 $^{^{\}ddagger} Electronic address: katarzyna.lewandowska@gumed.edu.pl$

above have also been considered. The examples are the equations with Caputo–Fabrizio fractional time derivative [34], with tempered Caputo derivative [35], and with Hilfer fractional derivative with respect to another function [36].

We consider a Cattaneo-type subdiffusion equation (CTSE) which is defined here as an equation that differs from the ordinary subdiffusion one by an additional operator acting on the time variable. The operator is generated by the constitutive equation describing the relation between the probability density flux and the gradient of Green's function. Combining this equation with the continuity equation the CTSE is obtained. As we show in Sec. VII, under certain assumptions, the equation can also be derived within the ordinary continuous time random walk model, in which the distribution of the waiting time for a molecule to jump is controlled by the additional operator. The Cattaneo-type subdiffusion equation takes into account a process, called the additional Cattaneo-type process (ACTP), that change ordinary subdiffusion. An example is diffusion of antibiotics in bacterial biofilms. In a dense biofilm subdiffusion of antibiotic molecules may occur. However, bacteria can activate defense mechanisms against the effects of antibiotics. Some of them additionally slow down subdiffusion or temporarily trap antibiotic molecules [37, 38].

In the case of passive subdiffusion, the solutions to the Cattaneo equation and the ordinary subdiffusion one are usually not much different from each other. Then, the latter equation, which appears to be easier to solve, is used to describe diffusion. However, there are processes that both equations provide qualitatively different results even for small τ . Examples of this are diffusive and subdiffusive impedance [39–42]. When considering subdiffusion arises whether such an equation gives a significant difference compared to the use of the ordinary subdiffusion equation. Therefore, we consider functions describing how the ACTP affects ordinary subdiffusion.

The general subdiffusion equation controlled by a memory kernel (MK) (equivalent to Eqs. (13)–(15) presented later), which is the basis for deriving the Cattaneo-type equation, has been considered to derive various types of anomalous diffusion equations. Examples are equations generated by distributed order MK [43–47], power–law MK [44], and truncated power law MK [43]. The equation with MK has been used to derive multi-fractional diffusion equations [26, 45–47]. Generalized diffusion equations with MK can also be derived by means of the subordination method [48].

The type of diffusion is often defined by the time evolution of the mean square displacement (MSD) of a diffusing particle σ^2 . When MSD is a power function of time, $\sigma^2(t) \sim t^{\alpha}$, $\alpha > 1$ is for superdiffusion, $\alpha = 1$ for normal diffusion, and $0 < \alpha < 1$ is for subdiffusion, when MSD is expressed by a slowly varying function, ultraslow diffusion (slow subdiffusion) occurs. In superdiffusion, a diffusing molecule can perform anomalously long jumps with a relatively high probability, while in subdiffusion the waiting time for a molecule to jump can be anomalously long. However, MSD alone does not clearly determine the type of diffusion [49, 50]. As shown in Ref. [50], the appropriate combination of superdiffusion and subdiffusion effects leads to a process in which $\sigma^2(t) \sim t$, but such process is certainly not normal diffusion. In addition to MSD, we also consider the first passage time distribution F(t;x) of a particle through an arbitrarily chosen point x. When diffusion in a subdiffusive medium is described by the Cattaneo-type equation, for short time there is $\sigma^2(t) \sim t^{\nu}$ with $\nu > 1$ [16], $\sigma^2(t) \sim t^{\alpha}$ with $\alpha \in (0,1)$ is achieved in the long-time limit. The appearance of a term with an exponent greater than 1 is typical for superdiffusion. However, it is difficult to expect that superdiffusion could be observed in a subdiffusive medium which properties do not change over time. The question arises whether the type of diffusion is determined only by the properties of the medium and the diffusing substance and if it can be verified by an experiment or whether it may depend on the model describing diffusion. Leaving aside the interpretative ambiguities, an equation whose Green's function provides the MSD with the exponent $\alpha \in (0,1)$ when $t \to \infty$ is referred to as a subdiffusion equation.

We consider subdiffusion in a one-dimensional homogeneous system with constant parameters. We pay attention to the interpretation of ACTP and assesses how this process affects subdiffusion. In Sec. II the subdiffusion equation generated by the generalized flux equation controlled by a MK β is derived. In Sec. III we consider functions describing subdiffusion. The explicit forms of the functions are given in terms of the Laplace transform. In Sec. IV the function β linear with respect to the parameter τ is considered. This function generates a Cattaneo-type subdiffusion equation. The CTSEs with the additional Caputo fractional time derivative of the order independent of the subdiffusion exponent and with the additional operator generated by a slowly varying function are studied in Secs. V and VI, respectively. We show how to derive the CTSE within the ordinary continuous time random walk model in Sec. VII. Final remarks are presented in Sec. VIII.

II. SUBDIFFUSION EQUATION GENERATED BY GENERALIZED FLUX EQUATION

The subdiffusion equation can be obtained phenomenologically by a combination of the continuity equation

$$\frac{\partial P(x,t)}{\partial t} = -\frac{\partial J(x,t)}{\partial x},\tag{1}$$

and the constitutive equation that defines the flux J. For ordinary subdiffusion the flux is defined as

$$J(x,t) = -D\frac{^{RL}\partial^{1-\alpha}}{\partial t^{1-\alpha}}\frac{\partial P(x,t)}{\partial x},$$
(2)

where $\alpha \in (0, 1)$, with the Riemann–Liouville fractional derivative

$$\frac{RL}{dt^{\nu}} \frac{d^{\nu} f(t)}{dt^{\nu}} = \frac{1}{\Gamma(n-\nu)} \frac{d^n}{dt^n} \int_0^t (t-u)^{n-\nu-1} f(u) du, \quad (3)$$

n is a natural number, $n = [\nu] + 1$ when $\nu \notin \mathcal{N}$ and $n = \nu \in \mathcal{N}$. Throughout this paper we assume that the initial particle position is $x_0 = 0$. Combining Eqs. (1) and (2) one gets

$$\frac{\partial P(x,t)}{\partial t} = D \frac{{}^{RL} \partial^{1-\alpha}}{\partial t^{1-\alpha}} \frac{\partial^2 P(x,t)}{\partial x^2}.$$
 (4)

When deriving a diffusion equation, it is convenient to use the Laplace transform

$$\mathcal{L}[f(t)](s) = \int_0^\infty e^{-st} f(t) dt \equiv \hat{f}(s).$$
 (5)

The Laplace transforms of derivatives are

$$\mathcal{L}\left[\frac{^{RL}d_{g}^{\alpha}f(t)}{dt^{\alpha}}\right](s) = s^{\alpha}\mathcal{L}\left[f(t)\right](s), \ \alpha \in (0,1), \quad (6)$$

$$\mathcal{L}\left[\frac{d^{n}f(t)}{dt^{n}}\right] = s^{n}\mathcal{L}[f(t)](s) - \sum_{i=0}^{n-1} s^{n-1-i}f^{(i)}(0), \quad (7)$$

where $f^{(i)}(t) = d^i f(t)/dt^i$, *i* and *n* are natural numbers. The Laplace transforms of Eqs. (1), (2), and (4) are, respectively,

$$s\hat{P}(x,s) - P(x,0) = -\frac{\partial J(x,s)}{\partial x},$$
(8)

$$\hat{J}(x,s) = -Ds^{1-\alpha} \frac{\partial \hat{P}(x,s)}{\partial x},$$
(9)

$$s\hat{P}(x,s) - P(x,0) = Ds^{1-\alpha} \frac{\partial^2 \hat{P}(x,s)}{\partial x^2}.$$
 (10)

The assumption under which the Cattaneo diffusion equation was derived is that the flux is delayed with respect to the concentration gradient. However, in the case of subdiffusion, the relation between the flux and the concentration gradient can be assumed in many ways, see Refs. [16, 17]. We assume that

$$\hat{\beta}(s;\tau)\hat{J}(x,s) = -Ds^{1-\alpha}\frac{\partial \hat{P}(x,s)}{\partial x},\qquad(11)$$

 β is a memory kernel which depends on the parameter τ . We also assume that for $\tau = 0$ we get ordinary subdiffusion equation, then

$$\hat{\beta}(s,0) = 1. \tag{12}$$

Subdiffusion is a non–Markovian process with a "long memory" controlled by the parameter $\alpha \in (0, 1)$. Within the ordinary continuous time random walk (CTRW) model, the probability density of waiting time for a molecule to jump ψ has a heavy tail, $\psi(t) \sim 1/t^{1+\alpha}$, the average value of the time is then infinite. This effect is caused by an internal structure of the medium, which makes it difficult for molecules to move. The model leads to the ordinary subdiffusion equation Eq. (4). In some cases an additional process retarding the flux with respect to the concentration gradient can be included in the model. In our considerations, such additional process is described by the function β , which may be independent of the parameter α . Stochastic interpretation of the process is discussed in Sec. VII. Eqs. (8) and (11)provide

$$\hat{\beta}(s;\tau) \left[s\hat{P}(x,s) - P(x,0) \right] = Ds^{1-\alpha} \frac{\partial^2 P(x,s)}{\partial x^2}.$$
 (13)

In the time domain Eq. (13) reads

$$\mathcal{B}_{t;\tau}[P(x,t)] = D \frac{RL\partial^{1-\alpha}}{\partial t^{1-\alpha}} \frac{\partial^2 P(x,t)}{\partial x^2}, \qquad (14)$$

where

$$\mathcal{B}_{t;\tau}[f(t)] = \int_0^t \beta(t - t'; \tau) f^{(1)}(t') dt'.$$
 (15)

Constraints on the function β are presented in the following sections. For $\hat{\beta}(s;\tau) = 1 + \tau s$ and $\alpha = 1$ we get the well-known Cattaneo diffusion equation,

$$\tau \frac{\partial^2 P(x,t)}{\partial t^2} + \frac{\partial P(x,t)}{\partial t} = D \frac{\partial^2 P(x,t)}{\partial x^2}.$$
 (16)

III. FUNCTIONS DESCRIBING DIFFUSION

The Green's function is the solution to diffusion equation with the boundary conditions

$$P(\pm\infty, t) = 0, \tag{17}$$

and the initial condition

$$P(x,0) = \delta(x), \tag{18}$$

where δ is the delta-Dirac function. The above initial condition is sufficient to determine the Laplace transform of a time derivative of at most first order. When computing the transform of other operators additional initial conditions may be needed. We assume that the only non-zero initial condition is Eq. (18).

In terms of the Laplace transform the Green's function of Eq. (14) is

$$\hat{P}(x,s) = \frac{\sqrt{\hat{\beta}(s;\tau)}}{2\sqrt{D}s^{1-\alpha/2}} e^{-\frac{s^{\alpha/2}|x|\sqrt{\hat{\beta}(s;\tau)}}{\sqrt{D}}}.$$
 (19)

Eq. (19) takes the form of Eq. (13) derived in Ref. [28] (putting $\hat{M}_{n,\gamma}(s) = s^{\alpha} \hat{\beta}(s;\tau)$ and B = D in the latter equation we get Eq. (19)). Since P(x,t) is a probability density, this function must be normalized and non-negative. Integrating Eq. (19) we get $\int_{-\infty}^{\infty} \hat{P}(x,s) dx = 1/s$, thus the normalization condition is met. The challenge is to determine conditions for the function β that ensure the non-negativity of the function P. The non-negativity property can be checked for various β using analytical or numerical methods. However, there are some general conditions imposed on $\hat{\beta}$ ensuring non-negativity of P. Guided by the method presented in Ref. [28], we find that P is certainly non-negative when $\hat{\beta}$ is a complete Bernstein function (CBF). The proof of this statement uses conclusion from the Bernstein theorem that the function P(x,t) is non-negative when $\hat{P}(x,s)$ is a completely monotone function (\mathcal{CMF}) , i.e. $(-1)^n \partial^n \hat{P}(x,s) / \partial s^n \ge 0$ for $n = 0, 1, 2, \dots$ According to the properties of \mathcal{CMF} and \mathcal{CBF} , presented in Refs. [28, 51] and in a shortened version in Appendix A in this paper, $\hat{P} \in CMF$ when $\hat{\beta} \in CBF$.

The Green's function Eq. (19) is symmetrical, P(x,t) = P(-x,t), this property is used in deriving further equations. On the basis of the Green's function, other functions characterizing diffusion processes are determined. A frequently used function is the temporal evolution of the mean square displacement σ^2 of diffusing particle,

$$\sigma^2(t) = 2 \int_0^\infty x^2 P(x,t) dx.$$
⁽²⁰⁾

In terms of the Laplace transform we get

$$\hat{\sigma^2}(s) = \frac{2D}{s^{1+\alpha}\hat{\beta}(s;\tau)}.$$
(21)

Another function that characterizes diffusion is the first passage time distribution F. Assuming x > 0, the distribution of time of the first particle passing the point xcan be calculated by means of the formula

$$F(t;x) = -\frac{\partial}{\partial t} \int_{-x}^{x} P(x',t) dx'.$$
 (22)

In terms of the Laplace transform the above equation reads

$$\hat{F}(s;x) = 1 - s \int_{-x}^{x} \hat{P}(x',s) dx'.$$
 (23)

Putting Eq. (19) to Eq. (23) we get

$$\hat{F}(s;x) = e^{-\frac{xs^{\alpha/2}\sqrt{\hat{\beta}(s;\tau)}}{\sqrt{D}}}.$$
(24)

The probability flux at x is defined as

$$J(x,t) = -\frac{\partial}{\partial t} \int_{-\infty}^{x} P(x',t) dx'.$$
 (25)

The Laplace transform of Eq. (25) is

$$\hat{J}(x,s) = 1 - s \int_{-\infty}^{x} \hat{P}(x',s) dx'.$$
 (26)

For x > 0 we obtain

$$\hat{J}(x,s) = \frac{1}{2} e^{-\frac{xs^{\alpha/2}\sqrt{\hat{\beta}(s;\tau)}}{\sqrt{D}}}.$$
 (27)

The equations (24) and (27) provide the relation

$$J(x,t) = \frac{1}{2}F(t;x), \ x > 0.$$
 (28)

For x < 0 the flux is negative, then J(x,t) = -F(t;x)/2.

The question arises whether the Cattaneo subdiffusion equation brings a new quality in comparison with the ordinary subdiffusion equation. We compare the functions F and σ^2 derived for $\tau = 0$ and for $\tau \neq 0$. The functions F_R and σ_R^2 show the time evolution of the relative change of first passage time distribution and MSD, respectively,

$$F_R(t;x) = \frac{F_{\tau=0}(t;x) - F(t;x)}{F_{\tau=0}(t;x)},$$
(29)

$$\sigma_R^2(t) = \frac{\sigma_{\tau=0}^2(t) - \sigma^2(t)}{\sigma_{\tau=0}^2(t)},$$
(30)

in Eq. (29) x is a parameter. Additionally, we consider the relative change of the Green's function

$$P_R(x,t) = \frac{P_{\tau=0}(x,t) - P(x,t)}{P_{\tau=0}(x,t)},$$
(31)

here x is a random variable, t is a parameter.

IV. CATTANEO-TYPE SUBDIFFUSION EQUATION

Let us assume that

$$\hat{\beta}(s;\tau) = 1 + \tau \hat{\gamma}(s). \tag{32}$$

Assuming that $\hat{\gamma}(s) \in CBF$ we have $\hat{\beta} \in CBF$, due to property 3 in Appendix A. As we argued earlier, the last relation ensures the non-negativity of the function P. Combining Eqs. (13) and (32) we obtain

$$(1 + \tau \hat{\gamma}(s)) \left[s \hat{P}(x,s) - P(x,0) \right]$$
(33)
$$= Ds^{1-\alpha} \frac{\partial^2 \hat{P}(x,s)}{\partial x^2}.$$

The inverse Laplace transform of Eq. (33) provides the Cattaneo-type subdiffusion equation

$$\tau \mathcal{D}_t[P(x,t)] + \frac{\partial P(x,t)}{\partial t} = D \frac{^{RL}\partial^{1-\alpha}}{\partial t^{1-\alpha}} \frac{\partial^2 P(x,t)}{\partial x^2}, \quad (34)$$

where

$$\mathcal{D}_t[f(t)] = \int_0^t \gamma(t - t') f^{(1)}(t') dt'.$$
 (35)

The function γ is determined by the ACTP. We note that for some γ the operator \mathcal{D}_t can be interpreted as a fractional time derivative.

The Laplace transforms of Green's function and first passage time distribution are

$$\hat{P}(x,s) = \frac{\sqrt{1+\tau\hat{\gamma}(s)}}{2\sqrt{D}s^{1-\alpha/2}} e^{-\frac{s^{\alpha/2}|x|\sqrt{1+\tau\hat{\gamma}(s)}}{\sqrt{D}}},\qquad(36)$$

$$\hat{F}(s;x) = e^{-\frac{xs^{\alpha/2}\sqrt{1+\tau\hat{\gamma}(s)}}{\sqrt{D}}}.$$
(37)

Assuming $\tau \hat{\gamma}(s) < 1$, the above functions can be presented as power series with respect to τ ,

$$\hat{P}(x,s) = \frac{1}{2\sqrt{D}s^{1-\alpha/2}} e^{-\frac{|x|s^{\alpha/2}}{\sqrt{D}}} \sum_{i=0}^{\infty} \tau^{i} a_{i}(x,s) \hat{\gamma}^{i}(s), \quad (38)$$

$$\hat{F}(s;x) = e^{-\frac{|x|s^{\alpha/2}}{\sqrt{D}}} \sum_{i=0}^{\infty} \tau^{i} b_{i}(x,s) \hat{\gamma}^{i}(s).$$
(39)

The functions $a_i(x,s)$ and $b_i(x,s)$ are determined applying the following series $\sqrt{1+u} = 1 + u/2 + \sum_{\substack{n=2\\n=0}}^{\infty} (-1)^{n-1} (2n-1)!! u^n / 2^n n!$, |u| < 1, and $e^{-u} = \sum_{\substack{n=0\\n=0}}^{\infty} (-1)^n u^n / n!$. In the following, we use the approximation of \hat{P} and \hat{F} including four leading terms in the series. For $0 \le i \le 3$ we have

$$a_0(x,s) = 1, \ a_1(x,s) = \frac{1}{2} \left(1 - \frac{|x|s^{\alpha/2}}{\sqrt{D}} \right),$$
 (40)

$$a_2(x,s) = \frac{1}{8} \Big(-1 - \frac{|x|s^{\alpha/2}}{\sqrt{D}} + \frac{|x|^2 s^{\alpha}}{D} \Big), \qquad (41)$$

$$a_3(x,s) = \frac{1}{16} \left(1 + \frac{|x|s^{\alpha/2}}{\sqrt{D}} - \frac{|x|^{3/2}s^{3\alpha/2}}{3D^{3/2}} \right), \quad (42)$$

$$b_0(x,s) = 1, \ b_1(x,s) = -\frac{|x|s^{\alpha/2}}{2\sqrt{D}},$$
 (43)

$$b_2(x,s) = \frac{|x|s^{\alpha/2}}{8\sqrt{D}} \left(1 + \frac{|x|s^{\alpha/2}}{\sqrt{D}}\right), \quad (44)$$

$$b_3(x,s) = \frac{-|x|s^{\alpha/2}}{16\sqrt{D}} \left(12 + \frac{|x|s^{\alpha/2}}{\sqrt{D}} + \frac{|x|^2s^{\alpha}}{3D}\right).$$
 (45)

From Eqs. (21), and (32) we get

$$\hat{\sigma}^2(s) = \frac{2D}{s^{1+\alpha}(1+\tau\hat{\gamma}(s))}.$$
(46)

The series representation of the above function is

$$\hat{\sigma^2}(s) = \frac{2D}{\Gamma(1+\alpha)s^{1+\alpha}} \sum_{i=0}^{\infty} \tau^i (-\hat{\gamma}(s))^i. \quad (47)$$
CTSE WITH THE ADDITIONAL CAPUTO
FRACTIONAL TIME DERIVATIVE

Let

v.

$$\hat{\gamma}(s) = s^{\kappa}, \ 0 < \kappa \le 1.$$
(48)

In terms of the Laplace transform we get

$$(\tau s^{\kappa} + 1) \left[s \hat{P}(x,s) - P(x,0) \right] = D s^{1-\alpha} \frac{\partial^2 \hat{P}(x,s)}{\partial x^2}.$$
 (49)

Using the relation

$$\mathcal{L}\left[\frac{^{C}\partial^{\alpha}f(t)}{\partial t^{\alpha}}\right](s) = s^{\alpha}\hat{f}(s) - \sum_{i=0}^{n-1}s^{\alpha-1-i}f^{(i)}(0), \quad (50)$$

where

$$\frac{^{C}d^{\alpha}f(t)}{dt^{\alpha}} = \frac{1}{\Gamma(n-\alpha)} \int_{0}^{t} (t-t')^{n-\alpha-1} f^{(n)}(t') dt' \quad (51)$$

is the Caputo fractional derivative (the natural number n is defined as under Eq. (3)), we get

$$\tau \frac{C\partial^{1+\kappa} P(x,t)}{\partial t^{1+\kappa}} + \frac{\partial P(x,t)}{\partial t} = D \frac{RL\partial^{1-\alpha}}{\partial t^{1-\alpha}} \frac{\partial^2 P(x,t)}{\partial x^2},$$
(52)

the initial conditions are $P(x, 0) = \delta(x)$ and $P^{(1)}(x, 0) = 0$.

The following formula plays a key role in calculating the inverse Laplace transforms [52]

$$\mathcal{L}^{-1}\left[s^{\nu}\mathrm{e}^{-as^{\beta}}\right](t) \equiv f_{\nu,\beta}(t;a)$$

$$= \frac{1}{t^{1+\nu}} \sum_{j=0}^{\infty} \frac{1}{j!\Gamma(-\nu-\beta j)} \left(-\frac{a}{t^{\beta}}\right)^{j},$$
(53)

 $a, \beta > 0, f_{\nu,\beta}$ is a special case of the Fox H–function (see Refs. [10, 52] and the references cited therein),

$$f_{\nu,\beta}(t;a) = \frac{1}{\beta a^{(1+\nu)/\beta}} H_{11}^{10} \left(\frac{a^{1/\beta}}{t} \middle| \begin{array}{c} 1 & 1 \\ (1+\nu)/\beta & 1/\beta \end{array} \right) \ .$$

From Eqs. (38)-(45) and Eq. (53) we get

$$P(x,t) = \frac{1}{2\sqrt{D}} \left[\xi_{-1+\alpha/2}(|x|,t) + \frac{\tau}{2} \left(\xi_{-1+\kappa+\alpha/2}(|x|,t) - \frac{|x|}{\sqrt{D}} \xi_{-1+\kappa+\alpha}(|x|,t) \right) + \frac{\tau^2}{8} \left(-\xi_{-1+2\kappa+\alpha/2}(|x|,t) - \frac{|x|}{\sqrt{D}} \xi_{-1+2\kappa+\alpha}(|x|,t) + \frac{|x|^2}{D} \xi_{-1+2\kappa+3\alpha/2}(|x|,t) \right) + \frac{\tau^3}{16} \left(\xi_{-1+3\kappa+\alpha/2}(|x|,t) + \frac{|x|}{\sqrt{D}} \xi_{-1+3\kappa+\alpha}(|x|,t) - \frac{|x|^3}{3D^{3/2}} \xi_{-1+3\kappa+2\alpha}(|x|,t) \right) \right],$$
(54)

$$F(t;x) = \xi_0(|x|,t) - \frac{\tau|x|}{2\sqrt{D}}\xi_{\kappa+\alpha/2}(|x|,t) + \frac{\tau^2|x|}{8\sqrt{D}}\left(\xi_{2\kappa+\alpha/2}(|x|,t) + \frac{|x|}{\sqrt{D}}\xi_{2\kappa+\alpha}(|x|,t)\right) - \frac{\tau^3|x|}{16\sqrt{D}}\left(12\xi_{3\kappa+\alpha/2}(|x|,t) + \frac{|x|}{\sqrt{D}}\xi_{3\kappa+\alpha}(|x|,t) + \frac{|x|^2}{3D}\xi_{3\kappa+3\alpha/2}(|x|,t)\right),$$
(55)

where

$$\xi_{\nu}(x,t) \equiv f_{\nu,\alpha/2}\left(t;\frac{x}{\sqrt{D}}\right).$$
(56)

A special case of this function for $\nu = -1 + \alpha/2$ is called the Mainardi function.

Assuming $\tau s^{\kappa} < 1$ we obtain

$$\hat{\sigma^2}(s) = \frac{2D}{s^{1+\alpha}} \sum_{i=0}^{\infty} (-\tau)^i s^{\kappa i}.$$
 (57)

It is not possible to calculate the inverse Laplace transform of the series in the above equation term by term. Therefore, we calculate the transform of $e^{-as^{\mu}}\hat{\sigma}^2(s)$, $a, \mu > 0$, using Eq. (53), next the limit of the obtained function when $a \to 0^+$ is calculated; the result is independent of the parameter μ . We obtain

$$\sigma^2(t) = 2Dt^{\alpha} \sum_{i=0}^{\infty} \frac{(-\tau)^i}{\Gamma(1+\alpha-i\kappa)t^{i\kappa}}.$$
 (58)

In the long time limit the functions Eqs. (55) and (58) generate the following relations

$$F_R(t \to \infty; x) = A_\kappa \frac{\tau}{t^\kappa},\tag{59}$$

$$\sigma_R^2(t \to \infty) = B_\kappa \frac{\tau}{t^\kappa},\tag{60}$$

where $A_{\kappa} = -\Gamma(-\alpha/2)/[2\Gamma(-\kappa - \alpha/2)]$ and $B_{\kappa} = \Gamma(1 + \alpha)/\Gamma(1 + \alpha - \kappa)$.

The figures 1–3 show the plots of the relative functions σ_R^2 , F_R , and P_R , respectively. Values of variables and parameters are given in arbitrarily chosen units. When ACTP has a large impact on subdiffusion the relative functions take large values and when the impact is small the functions are close to zero. Figs. 1 and 2 suggest that the ACTP disappears over time, the main influence



FIG. 1: Time evolution of the relative MSD for the parameters given in the legend, $\tau = 0.1$ and D = 5.

on the functions is the parameter κ . In Fig. 3 we see that the ACTP increases with moving away from the initial position of a particle. This effect causes that the probability of finding the particle far from its initial position is much lower when the process is described by the Cattaneo-type subdiffusion equation.





FIG. 2: Time evolution of the relative first passage time for the parameters given in the legend, x = 10, $\tau = 0.1$, and D = 5.

VI. CTSE WITH AN ADDITIONAL OPERATOR GENERATED BY A SLOWLY VARYING FUNCTION

A slowly varying function $\mathcal{R}(t)$ at infinity fulfils the relation

$$\frac{\mathcal{R}(at)}{\mathcal{R}(t)} \to 1 \tag{61}$$

when $t \to \infty$ for any a > 0. Functions that have finite limits when $t \to \infty$ and logarithmic functions are examples of slowly varying functions.

In the following, we use the strong Tauberian theorem to determine the inverse Laplace transform in the longtime limit [53]: if $\phi(t) \geq 0$, $\phi(t)$ is ultimately monotonic like $t \to \infty$, \mathcal{R} is slowly varying at infinity, and $0 < \rho < \infty$, then each of the relations

$$\hat{\phi}(s) \approx \frac{\mathcal{R}(1/s)}{s^{\rho}}$$
 (62)

as $s \to 0$ and

$$\phi(t) \approx \frac{\mathcal{R}(t)}{\Gamma(\rho)t^{1-\rho}} \tag{63}$$

as $t \to \infty$ implies the other.

Let $\hat{\gamma}(s) \equiv \mathcal{R}(1/s)$ be a slowly varying function. When

FIG. 3: The relative Green's function P_R for α and κ given in the legend, t = 20, $\tau = 0.1$, and D = 10.

 $s \to 0$, the Green's function Eq. (36) is approximated as

$$\hat{P}(x,s) = \frac{\sqrt{1 + \tau \mathcal{R}(1/s)}}{2\sqrt{D}s^{1-\alpha/2}}$$

$$\times \left[1 - \frac{s^{\alpha/2}|x|\sqrt{1 + \tau \mathcal{R}(1/s)}}{\sqrt{D}}\right].$$
(64)

Using the strong Tauberian theorem we get in the long time limit

$$P(x,t) = \frac{\sqrt{1 + \tau \mathcal{R}(t)}}{2\sqrt{D}\Gamma(1 - \alpha/2)t^{\alpha/2}}$$

$$\times \left[1 - \frac{\Gamma(1 - \alpha/2)\sqrt{1 + \tau \mathcal{R}(t)}|x|}{\sqrt{D}\Gamma(1 - \alpha)t^{\alpha/2}}\right].$$
(65)

Putting the above equation to Eq. (22) we get

$$F(t \to \infty; x) = \frac{|x|}{2\sqrt{D}\Gamma(1 - \alpha/2)\sqrt{1 + \tau \mathcal{R}(t)}t^{\alpha/2}} \quad (66)$$
$$\times \left[\frac{1 + \tau \mathcal{R}(t)}{t} - \tau \mathcal{R}^{(1)}(t)\right].$$

Eqs. (29) and (66) provide

$$F_R(t \to \infty; x) = 1 - \sqrt{1 + \tau \mathcal{R}(t)}$$

$$+ \frac{\tau t \mathcal{R}^{(1)}(t)}{\sqrt{1 + \tau \mathcal{R}(t)}}.$$
(67)

From Eqs. (30), (46), and the strong Tauberian theorem we obtain

$$\sigma^{2}(t \to \infty) = \frac{2Dt^{\alpha}}{\Gamma(1+\alpha)(1+\tau \mathcal{R}(t))}, \qquad (68)$$

and

$$\sigma_R^2(t \to \infty) = \frac{\tau \mathcal{R}(t)}{1 + \tau \mathcal{R}(t)}.$$
 (69)

As an example we consider the CTSE with

$$\hat{\gamma}(s) \equiv \mathcal{R}(1/s) = \frac{\log(1 + a\sqrt{s})}{\sqrt{s}},\tag{70}$$

a > 0. The above function is a slowly varying function as well as a complete Bernstein function, see Ch. 15.4 in Ref. [51]. There is

$$\mathcal{R}(t) = \sqrt{t} \log\left(1 + \frac{a}{\sqrt{t}}\right). \tag{71}$$

In the limit of long time we get

$$F_R(t \to \infty; x) = 1 - \sqrt{1 + \tau a}, \qquad (72)$$

$$\sigma_R^2(t \to \infty) = \frac{\tau a}{1 + \tau a}.$$
(73)

Thus, the ACTP effect is visible for a long time.

VII. CTSE DERIVED FROM THE ORDINARY CTRW MODEL

In our considerations, we derived the Cattaneo-type equation assuming that the flux is not generated at the same time as the concentration gradient. The time shift between the flux and the concentration gradient is controlled by the function β . However, there are other models that provide the Cattaneo equation. The equation has been derived within the continuous time persistent random walk (CTPRW) model, see Refs. [28, 54, 55] and the references cited therein. Unlike in the ordinary continuous time random walk (CTRW) model [56], in CT-PRW model the probability of choosing the direction of a molecule jump depends on the direction of its previous jump. In this section we show that, under certain conditions imposed on functions β and γ , the Cattaneo-type equation can be derived by means of the ordinary CTRW model with a properly defined distribution of waiting time for a molecule to jump.

The ordinary CTRW model provides the following equation given in terms of the Laplace transform, see Appendix B,

$$s\hat{P}(x,s) - P(x,0) = \frac{\epsilon^2 s\hat{\psi}(s)}{2[1-\hat{\psi}(s)]} \frac{\partial^2 \hat{P}(x,s)}{\partial x^2}, \qquad (74)$$

where ϵ is the mean value of a single jump length. Comparing Eqs. (13) and (74) we obtain

$$\hat{\psi}(s) = \frac{1}{1 + \lambda s^{\alpha} \hat{\beta}(s;\tau)},\tag{75}$$

where $\lambda = \epsilon^2/2D$. The parameter describing subdiffusion is D, ϵ is a scale-dependent parameter. We further assume that ϵ is small, which provides $\lambda \ll 1$. Guided by the considerations presented in Ref. [28] we conclude that the function ψ Eq. (75) is non-negative if $s^{\alpha}\hat{\beta}(s;\tau) \in \mathcal{CBF}$. Due to the normalization condition, $\hat{\psi}(0) = 1$, we additionally assume that $s^{\alpha}\hat{\beta}(s;\tau) \to 0$ when $s \to 0$.

From Eqs. (32) and (75) we get

$$\hat{\psi}(s) = \frac{1}{1 + \lambda s^{\alpha} + \lambda \tau s^{\alpha} \hat{\gamma}(s)}.$$
(76)

The representation of the function ψ Eq. (76) as a power series with respect to τ is

$$\hat{\psi}(s) = \sum_{i=0}^{\infty} \tau^i \hat{\gamma}^i(s) \sum_{n=i}^{\infty} \binom{n}{i} (-\lambda)^n s^{n\alpha}.$$
 (77)

When $\lambda \ll 1$ we have

$$\hat{\psi}(s) \approx \hat{\psi}_{\tau=0}(s)\hat{\psi}_{\gamma}(s), \tag{78}$$

where

$$\hat{\psi}_{\tau=0}(s) = \frac{1}{1+\lambda s^{\alpha}},\tag{79}$$

$$\hat{\psi}_{\gamma}(s) = \frac{1}{1 + \lambda \tau s^{\alpha} \hat{\gamma}(s)}.$$
(80)

In the time domain Eq. (78) reads

$$\psi(t) \approx \int_0^t \psi_{\tau=0}(t-t')\psi_{\gamma}(t')dt'.$$
 (81)

The function $\psi_{\tau=0}$ is the distribution of the waiting time for a molecule to jump in a subdiffusive medium characterized by the parameters α and D, ψ_{γ} is the distribution of the waiting time when an additional mechanism disturbing the jump is turned on at the moment $t' \in (0, t)$.

As an example, we consider the ACTP process generated by $\hat{\gamma}(s) = s^{\mu}$. We assume $\mu \in (0, 1 - \alpha)$, then $s^{\alpha}\hat{\gamma}(s) \in CBF$ which ensures that also $\hat{\beta} \in CBF$. In this case we have

$$\hat{\psi}(s) = \frac{1}{1 + \lambda s^{\alpha} + \lambda \tau s^{\alpha + \mu}}.$$
(82)

Eqs. (74) and (82) provide the CTSE Eq. (52) for $\kappa = \mu < 1 - \alpha$. For sufficiently small s and/or λ the series expansion of the function $\hat{\psi}$ with respect to τ is

$$\hat{\psi}(s) = \sum_{n=0}^{\infty} \tau^n \sum_{i=n}^{\infty} (-\lambda)^i \binom{i}{n} s^{\alpha i + \mu n}.$$
(83)

Using the same procedure as for calculating Eq. (58), we get

$$\psi(t) = \sum_{n=0}^{\infty} \tau^n \sum_{i=n}^{\infty} \binom{i}{n} \frac{(-\lambda)^i}{\Gamma(-\alpha i - \mu n)t^{1+\alpha i + \mu n}}.$$
 (84)

In the limit of long time we have

$$\psi(t \to \infty) = \frac{-\lambda}{t^{1+\alpha}} \left[\frac{1}{\Gamma(-\alpha)} \right]$$

$$-\frac{\lambda}{\Gamma(-2\alpha)t^{\alpha}} + \frac{\tau}{\Gamma(-\alpha-\mu)t^{\mu}} .$$
(85)

Eq. (84) provides

$$\psi_{\tau=0}(t) = \sum_{i=1}^{\infty} \frac{(-\lambda)^i}{\Gamma(-\alpha i)t^{1+\alpha i}}.$$
(86)

Based on Eqs. (48) and (80) we get

$$\psi_{\gamma}(t) = \sum_{i=1}^{\infty} \frac{(-\lambda\tau)^i}{\Gamma(-(\alpha+\mu)i)t^{1+(\alpha+\mu)i}}.$$
(87)

For a short time (which corresponds to large s in the Laplace transform) we have $\psi_{\tau=0}(t) \sim 1/t^{1-\alpha}$ and $\psi_{\gamma}(t) \sim t^{\mu}/t^{1-\alpha}$, in the long time limit there is $\psi_{\tau=0}(t) \sim 1/t^{1+\alpha}$ and $\psi_{\gamma}(t) \sim 1/t^{1+\alpha+\mu}$. This suggests that for relatively short times the ACTP process is dominant over ordinary subdiffusion, the opposite is true for a long time.

VIII. FINAL REMARKS

In our considerations, the constitutive equation relating the flux and the concentration gradient is modified by the integral operator with kernel $\beta(t;\tau)$. The motivation for including the function β in the flux equation is to take into account the process of delaying the flux with respect to the concentration gradient; this process can be independent of the parameter α . In the time domain the equation reads

$$\int_{0}^{t} \beta(t-t';\tau) J(x,t') dt' = -D \frac{\partial^{1-\alpha}}{\partial t^{1-\alpha}} \frac{\partial P(x,t)}{\partial x}.$$
 (88)

Combining the above equation with the continuity equation we get the generalized subdiffusion equation. When $\hat{\beta}(s;\tau) = 1 + \tau \hat{\gamma}(s), \ \beta(t;\tau) = \delta(t) + \tau \gamma(t)$ in the time domain, we have the Cattaneo-type subdiffusion equation that differs from the ordinary subdiffusion one by an integro-differential operator with kernel $\gamma(t)$. When $\hat{\gamma}(s) = s^{\kappa}$ with $0 < \kappa < 1$ the operator is the Caputo fractional time derivative of the order $1 + \kappa$.

The functions σ_R^2 , F_R , and P_R can be interpreted as measures of the influence of ACTP on subdiffusion. For the process described by the CTSE with additional operator being the Caputo fractional time derivative there is $\sigma_R^2 \sim F_R \sim 1/t^{\kappa}$, $0 < \kappa < 1$, while for the equation with additional operator generated by a slowly varying function $\hat{\gamma}$ both functions σ_R^2 and F_R go to a nonzero constant when $t \to \infty$, see Eqs. (72) and (73). This example shows that the ACTP described by a slowly variable function may not vanish.

As we have mentioned in Sec. VII, another model providing the Cattaneo subdiffusion equation is CTPRW model [28, 54, 55]. Within this model, the directions of two consecutive molecule jumps are correlated, this may be motivated by an inertia of the molecule. The distribution of the waiting time for the molecule to jump can be chosen as Eq. (86). We have shown that the Cattaneo-type subdiffusion equation Eq. (34) can be derived within the ordinary CTRW model in which the distribution of the waiting time for a molecule to jump is the inverse Laplace transform of Eq. (76). This distribution has a heavy tail modified by ACTP, an example is shown in Eq. (85). It can be interpreted as a subdiffusion model which includes an additional process that affects the kinetics of subdiffusion. The molecule waits for a jump as in the ordinary subdiffusion process, and at any time an additional process can be activated that changes the probability density of molecule jump execution. In this model we do not consider the correlation of the successive molecule jumps.

An example of a process in which the CTSE effect can be created is subdiffusion (or normal diffusion when $\alpha = 1$) of antibiotics in a bacterial biofilm. The defense mechanisms of bacteria against the action of antibiotics can change the random walk of antibiotic molecules. It may be interesting to use the Cattaneo-type equation in modelling the spread of an epidemic. The first arrival of an infected object to some point may create new source of infection. Since ACTP changes the distribution of F, the use of Eqs. (4) and (52) (with possibly added reaction terms in both equations) in epidemic modelling may give qualitatively different results.

Appendix A: Completely monotone and complete Bernstein functions

A function $f: (0, \infty) \to \Re$ is a completely monotone function (\mathcal{CMF}) if f is of class C^{∞} and $(-1)^n f^{(n)}(u) \ge 0$ for $n = 0, 1, 2, \ldots$, where $f^{(n)}(u) = d^n f(u)/du^n$. The function f is a complete Bernstein function (\mathcal{CBF}) if f(u)/u is the Laplace transform of \mathcal{CMF} restricted to the positive semiaxis. The Brenstein theorem provides that $\hat{F}(s) \in \mathcal{CMF}$ if and only if $f(t) \ge 0$ for $t \ge 0$, where $\hat{F}(s) = \int_0^\infty \exp(-st)f(t)dt$. Some properties of the functions are [28, 51]:

- 1. $f, g \in CMF \Rightarrow f \cdot g \in CMF$,
- 2. $f \in CMF$ and $g \in CBF \Rightarrow f \circ g \equiv f(g) \in CMF$,
- 3. $f,g \in CBF \Rightarrow (f^{\alpha} + g^{\alpha})^{1/\alpha} \in CBF, \alpha \in [-1,1]/\{0\},$

4.
$$f, g \in CBF \Rightarrow f^{\alpha} \cdot g^{\beta} \in CBF, \ \alpha, \beta \in (0, 1), \ \alpha + \beta \leq 1,$$

5. $f(u) \in CBF(f(u) \neq 0) \Leftrightarrow u/f(u) \in CBF,$
6. $f(u) = \exp(-au) \in CMF, \ a > 0,$
7. $f(u) = u^{\mu} \in CMF \ when \ \mu \leq 0,$
8. $f(u) = u^{\mu} \in CBF \ when \ \mu \in [0, 1].$

Appendix B: Derivation of Eq. (74)

Let us suppose the probability density of the jump length Δx reads

$$\xi(\Delta x) = [\delta(\Delta x - \epsilon) + \delta(\Delta x + \epsilon)]/2, \qquad (89)$$

 $\epsilon < \infty$. In terms of the Fourier transform $f(k) \equiv \mathcal{F}[f(u)](k) = \int_{-\infty}^{\infty} \exp(iku)f(u)du$ we have $\tilde{\xi}(k) = \cos(k\epsilon)$; when $k\epsilon \to 0$, there is

$$\tilde{\xi}(k) = \rightarrow 1 - \frac{k^2 \epsilon^2}{2}.$$
 (90)

Combining Eq. (90) with the standard equation obtained within the ordinary CTRW model [56]

$$\hat{\tilde{P}}(k,s) = \frac{1 - \hat{\psi}(s)}{s} \frac{1}{1 - \tilde{\xi}(k)\hat{\psi}(s)},$$
(91)

- D. S. W. Lee, N. S. Wingreen, and C. P. Brangwynne, Chromatin mechanics dictates subdiffusion and coarsening dynamics of embedded condensates, Nat. Phys. 17, 531 (2021).
- [2] B. Bijeljic, P. Mostaghimi, and M. J. Blunt, Signature of non–Fickian solute transport in complex heterogeneous porous media, Phys. Rev. Lett. 107, 204502 (2011).
- [3] E. Barkai, Y. Garini, and R. Metlzer, Strange kinetics of single molecules in living cells, Phys. Today 65, 29 (2012).
- [4] T. Kosztołowicz, K. Dworecki, and S. Mrówczyński, How to measure subdiffusion parameters, Phys. Rev. Lett. 94, 170602 (2005).
- [5] T. Kosztołowicz and R. Metzler, Diffusion of antibiotics through a biofilm in the presence of diffusion and absorption barriers, Phys. Rev. E 102, 032408 (2020).
- [6] W. Wyss, The fractional diffusion equation, J. Math. Phys. 27, 2782 (1986).
- [7] R. Hilfer and L. Anton, Fractional master equations and fractal time random walks, Phys. Rev. E 51, R848 (1995).
- [8] A. Compte, Stochastic foundations of fractional dynamics, Phys. Rev. E 53, 4191 (1996).
- [9] R. Metzler, J. Klafter, and I. M. Sokolov, Anomalous transport in external fields: Continuous time random walks and fractional diffusion equations extended, Phys. Rev. E 58, 1621 (1998).
- [10] R. Metzler and J. Klafter, The random walk's guide to anomalous diffusion: a fractional dynamics approach,

we get

$$s\hat{\tilde{P}}(k,s) - 1 = -\frac{\epsilon^2 sk^2}{2[1 - \hat{\psi}(s)]}\hat{\tilde{P}}(k,s).$$
(92)

Using the formulas $\mathcal{F}^{-1}[1](x) = \delta(x) \equiv P(x,0)$ and $\mathcal{F}^{-1}[-k^2\hat{P}(k,s)](x) = \partial^2\hat{P}(x,s)/\partial x^2$ we get Eq. (74).

Another way to derive Eq. (74) is the particle random walk model based on the difference equation

$$P_{n+1}(x) = \frac{1}{2}P_n(x-\epsilon) + \frac{1}{2}P_n(x+\epsilon),$$
 (93)

where $P_n(x)$ is the probability density of finding the molecule at x after making n jumps. Combining Eq. (93) and the formulas $P(x,t) = \sum_{n=0}^{\infty} Q_n(t)P_n(x)$, where $Q_n(t)$ is the probability that a particle makes n jumps in the time interval (0,t) (in terms of the Laplace transform there is $\hat{Q}(s) = \hat{\psi}^n(s)(1-\hat{\psi}(s))/s)$ [56], and $\partial^2 P(x,t)/\partial x^2 \approx [P(x+\epsilon,t)+P(x-\epsilon,t)-2P(x,t)]/\epsilon^2$, we obtain Eq. (74) in the limit of small ϵ .

Phys. Rep. **339**, 1 (2000).

- [11] E. Barkai, R. Metzler, and J. Klafter, From continuous time random walks to the fractional Fokker-Planck equation, Phys. Rev. E 61, 132 (2000).
- [12] I. M. Sokolov, J. Klafter, and A. Blumen, Fractional kinetics, Phys. Today 55, 11, 48 (2002).
- [13] R. Klages, G. Radons, and I. M. Sokolov, Anomalous Transport: Foundations and Applications (Wiley, New York, 2008).
- [14] J. Klafter and I. M. Sokolov, First Step in Random Walks. From Tools to Applications (Oxford UP, New York, 2011).
- [15] C. Cattaneo, Sulla conduzione del calore. Attidel Seminario Matematico e Fisicodella Università di Modena 3, 83 (1948).
- [16] A. Compte and R. Metzler, The generalized Cattaneo equation for the description of anomalous transport processes, J. Phys. A: Math. Gen. 30, 7277 (1997).
- [17] E. Awad, On the time-fractional Cattaneo equation of distributed order, Physica A 518, 210 (2019).
- [18] G. Fernandez-Anaya, F. J. Valdes-Parada, and J. Alvarez-Ramirez, On generalized fractional Cattaneo's equations, Physica A **390**, 4198 (2011).
- [19] H. Qi and X. Jiang, Solutions of the space-time fractional Cattaneo equation, Physica A 390, 1876 (2011).
- [20] L. Liu, L. Zheng, and Y. Chen, Macroscopic and microscopic anomalous diffusion in comb model with fractional dual-phase-lag model, Appl. Math. Model. 62,

629 (2018).

- [21] S. D. Roscani, J. Bollati, and D. A. Tarzia, A new mathematical formulation for a phase change problem with a memory flux, Chaos Solit. Fract. 116, 340 (2018).
- [22] F. Alegria, V. Poblete, and J. C. Pozo, Nonlocal in-time telegraph equation and telegraph processes with random time, J. Differential Eqs. 347, 310 (2023).
- [23] Y. Luchko, F. Mainardi, and Y. Povstenko, Propagation speed of the maximum of the fundamental solution to the fractional diffusion-wave equation, Comput. Math. Appl. 66, 774 (2013).
- [24] E. Awad, T. Sandev, R. Metzler, and A. Chechkin, From continuous-time random walks to the fractional Jeffreys equation: Solution and properties, Int. J. Mass Heat Transf. 181, 121839 (2021).
- [25] Y. M. Hamada, Solution of a new model of fractional telegraph point reactor kinetics using differential transformation method, Appl. Math. Model. 78, 297 (2020).
- [26] R. Metzler and T. F. Nonnenmacher, Fractional diffusion, waiting-time distributions, and Cattaneo-type equations, Phys. Rev. E 57, 6409 (1998).
- [27] R. Metzler and A. Compte, Stochastic foundation of normal and anomalous Cattaneo-type transport, Physica A 268, 454 (1999).
- [28] K. Górska, A. Horzela, E. K. Lenzi, G. Pagnini, and T. Sandev, Generalized Cattaneo (telegrapher's) equations in modeling anomalous diffusion phenomena, Phys. Rev. E 102, 022128 (2020).
- [29] K. Górska, Integral decomposition for the solutions of the generalized Cattaneo equation, Phys. Rev. E 104, 024113 (2021).
- [30] T. Kosztołowicz, Cattaneo-type subdiffusion-reaction equation, Phys. Rev. E 90, 042151 (2014).
- [31] V. A. Vyawahare and P. S. V. Nataraj, Fractional-order modeling of neutron transport in a nuclear reactor, Appl. Math. Model. 37, 9747 (2013).
- [32] O. Nikan, Z. Avazzadeh, and J. A. Tenreiro Machado, Numerical approach for modeling fractional heat conduction in porous medium with the generalized Cattaneo model, Appl. Math. Model. **100**, 107 (2021).
- [33] M. Mozafarifard, D. Toghraie, and H. Sobhani, Numerical study of fast transient non-diffusive heat conduction in a porous medium composed of solid-glass spheres and air using fractional Cattaneo subdiffusion model, Int. Commun. Heat Mass Transf. **122**, 105192 (2021).
- [34] Z. Liu, A. Cheng, and X. Li, A second order Crank— Nicolson scheme for fractional Cattaneo equation based on new fractional derivative, Appl. Math. Comput. **311**, 361 (2017).
- [35] L. Beghin, R. Garra, F. Mainardi, and G. Pagnini, The tempered space-fractional Cattaneo equation, Probabilistic Engineering Mechanics 70, 103374 (2022).
- [36] N. Vieira, M. Ferreira, and M. M. Rodrigues, Timefractional telegraph equation with ψ-Hilfer derivatives, Chaos Solit. Fract. 162, 112276 (2022).
- [37] G. G. Anderson and G. A. O'Toole, Innate and induced resistance mechanisms of bacterial biofilms, Current Topics in Microbiology and Immunology **322**, 85 (2008).
- [38] T. F. C. Mah and G. A. O'Toole, Mechanisms of biofilm

resistance to antimicrobial agents, Trends in Microbiology 9(1), 34 (2001).

- [39] G. Barbero and J. Ross Macdonald, Transport process of ions in insulating media in the hyperbolic diffusion regime, Phys. Rev. E 81, 051503 (2010).
- [40] P. P. Kostrobij, I. I. Grygorchak, F. O. Ivaschyshyn, B. M. Markovych, O. V. Viznovych, and M. V. Tokarchuk, Mathematical modeling of subdiffusion impedance in multilayer nanostructures, Math. Model. Comput. 2, 154 (2015).
- [41] T. Kosztołowicz and K. D. Lewandowska, Hyperbolic subdiffusive impedance, J. Phys. A 42, 05500 (2009).
- [42] K. D. Lewandowska and T. Kosztołowicz, Appliaction of generalized Cattaneo equation to model subdiffusion impedance, Acta Phys. Polonica B 39, 1211 (2008).
- [43] T. Sandev, I. M. Sokolov, R. Metzler, and A. Chechkin, Beyond monofractional kinetics, Chaos Solit. Fract. 102, 210 (2017).
- [44] T. Sandev, A. Chechkin, H. Kantz, and R. Metzler, Diffusion and Fokker–Planck–Smoluchowski equations with generalized memory kernel, Fract. Cal. Appl. Anal. 18, 1006 (2015).
- [45] T. Sandev, N. Korabel, H. Kantz, I. M. Sokolov, and R. Metzler, Distributed order diffusion equations and multifractality: Models and solutions, Phys. Rev. E 92, 042117 (2015).
- [46] T. Sandev, R. Metzler, and A. Chechkin, From continuous time random walks to the generalized diffusion equation, Fract. Cal. Appl. Anal. 21, 10 (2018).
- [47] T. Sandev, Z. Tomovski, J. L. A. Dubbeldam, and A. Chechkin, Generalized diffusion–wave equation with memory kernel, J. Phys. A: Math. Theor. 52, 015201 (2019).
- [48] A. Chechkin and I. M. Sokolov, Relation between generalized diffusion equations and subodrination schemes, Phys. Rev. E 103, 032133 (2021).
- [49] Y. Meroz, I. M. Sokolov, and J. Klafter, Unequal twins: probability distributions do not determine everything, Phys. Rev. Lett. **107**, 260601 (2011).
- [50] B. Dybiec and E. Gudowska-Nowak, Discriminating between normal and anomalous random walks, Phys. Rev. E 80, 061122 (2009).
- [51] R.L. Schilling, R. Song, and Z. Vondraček, *Bernstein functions. Theory and applications* (De Gruyter, Berlin, 2010).
- [52] T. Kosztołowicz, From the solutions of diffusion equation to the solutions of subdiffusive one, J. Phys. A: Math. Gen. 37, 10779 (2004).
- [53] B. D. Hughes, Random Walk and Random Environments. Vol. I, Random Walks (Clarendon, Oxford, 1995).
- [54] J. Masoliver and K. Lindenberg, Continuous time persistent random walk: a review and some generalizations, Eur. Phys. J. B 90, 107 (2017).
- [55] J. Masoliver, Fractional telegrapher's equation from fractional persistent random walks, Phys. Rev. E 93, 052107 (2016).
- [56] E. W. Montroll and G. H. Weiss, Random walk on lattices. II, J. Math. Phys. 6, 167 (1965).