

FARMALINX

Dimop 375 EC

HERBICIDE

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply if rainfall is expected within 2 hours.
DO NOT spray when temperatures are higher than 25°C.
DO NOT apply to weeds or crop under stress due to, for e.g. very dry, very wet, nutrient deficient, frost or diseased conditions.
DO NOT use on ryegrass turf.

CROP	WEED	STATE	WEED STAGE	RATE L/ha	CRITICAL COMMENTS
Wheat, linseed, triticale, canola, safflower, rye, barley	Annual (Wimmera) Ryegrass	All States	2 to 4 leaf	1	Apply generally 3 to 6 weeks after sowing, when cereals usually 2 to 5 leaves. For barley, apply only when crop is 4 to 5 leaf stage. Add wetting agent at 0.25% a.i. to water in spray tank ie 250 mL WetDrop Wetter/100L. This is irrespective of volume of water/ha or rate of FARMALINX DIMOP 375 EC Herbicide. DO NOT add wetting agent if mixing with some other sprays (refer compatibility).
	Common Barbgrass	NSW only		1.25	
Lupins, peas	Annual Ryegrass	NSW, Vic, Tas, SA, WA only	2 to 4 leaf	1	
	Common Barbgrass	NSW only		1.25	
Wheat, linseed, triticale, safflower, rye, barley, canola	Wild Oats	All States	2 to 3 leaf	1.5	For barley, apply only when crop is 4 to 5 leaf stage. Avoid double spraying (overlap) at the maximum rate of 1.5 L/ha. In crops other than barley, use 2 L rate if most weeds 3 to 4 leaf and if spraying 4 to 6 weeks after sowing, especially in northern NSW and Qld. Add wetting agent as above.
			3 to 4 leaf	2 (not barley)	
Lupins, peas	Wild Oats	NSW, Vic, Tas, SA, WA only	2 to 3 leaf	1.5	Use 1.5 L rate only if growing conditions are very favourable and weeds are 2 to 3 leaf, 3 to 4 weeks after sowing, and cereals usually 2 to 3 leaf.
			3 to 4 leaf	2	
Pasture legume seed crops	Annual Ryegrass	All States	2 to 4 leaf	1	Add wetting agent as above. Ensure thorough coverage of weeds.
	Wild Oats			1.5 to 2	
Oilseed poppies	Wild Oats, Annual Ryegrass (refer critical comments)	Tas only	See critical comments	0.5 to 3	DO NOT add wetting agent. Apply in accordance with recommendations made by the Department of Agriculture or the poppy contracting company.

CROP	WEED	STATE	WEED STAGE	RATE L/ha	CRITICAL COMMENTS
Turf: kikuyu, hybrid couch cv. Tifdwarf	Crowsfoot Grass (<i>Eleusine indica</i>)	Qld, NSW, ACT only	See Critical Comments	1 L/ha or 100 mL in 100 L of water per 1000 m ²	Apply to weeds between the seedling and tillering stage for best results. Use lower rates for seedlings and higher rates for established grass weeds. A repeat spray may be needed in 2 to 4 weeks.
Queensland blue couch				1 to 1.5 L/ha or 100 to 150 mL in 100 L of water per 1000 m ²	
Common couch, buffalo, hybrid couch cv. Trifway, carpet grass, salt couch				1 to 2.5 L/ha or 100 to 250 mL in 100 L of water per 1000 m ²	Results are best when FARMALINX DIMOP 375 EC Herbicide is applied under good growing conditions and weeds have sufficient leaf area ie. 2 to 3 days after mowing.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIODS:

ALL CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR SEVEN WEEKS AFTER APPLICATION.
TURF: DO NOT ALLOW TREATED AREAS OR CLIPPINGS TO BE GRAZED BY OR FED TO LIVESTOCK.

GENERAL INSTRUCTIONS

Results are best under good growing conditions when weeds are actively growing. Application to weeds under stress eg Due to continuous severe frosts, dry and waterlogged conditions, nutrient deficiency or disease should be avoided. Wild Oats germinating after application will often be stunted and will not seriously compete with the crop. Application to crops with a covering of frost or dew has been successful but should be avoided if spraying cannot be completed.
If spraying cannot be completed spray mixtures of DIMOP 375 EC in clean water may be left overnight without significant loss of efficacy. The mixture should be well agitated before spraying re-commences. It is not recommended that this be general practice.

RESISTANT WEEDS WARNING

FARMALINX DIMOP 375 EC Herbicide is a member of the arylophenoxypropionate group of herbicides. FARMALINX DIMOP 375 EC Herbicide is an inhibitor of acetyl coA carboxylase. For weed resistance FARMALINX DIMOP 375 EC Herbicide is a Group A herbicide. Some naturally-occurring weed biotypes resistant to FARMALINX DIMOP 375 EC Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FARMALINX DIMOP 375 EC Herbicide or other Group A herbicides.
Since occurrence of resistant weeds is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of FARMALINX DIMOP 375 EC Herbicide to control resistant weeds.

GROUP A HERBICIDE

www.farmalinx.com

CONTINUED OVERLEAF

Additional information is listed in the safety data sheet (SDS). A safety data sheet for FARMALINX DIMOP 375 EC Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

SAFETY DATA SHEET

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water.

FIRST AID

Will irritate the eyes and skin. DO NOT inhale vapour. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

SAFETY DIRECTIONS

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

For reliable containers, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage. Rates. Wetting agent should be added to this mixture at the rate recommended on the FARMALINX DIMOP 375 EC Herbicide label. 5. Where phalaris is present, FARMALINX DIMOP 375 EC Herbicide and Puma[®]S may be applied together at the respective recommended rates. Wetting agent should be added to this mixture at the rate recommended on the FARMALINX DIMOP 375 EC Herbicide label. the Department of Agriculture or the poppy contracting company. 4. For use in oilseed poppies in Tasmania only: FARMALINX DIMOP 375 EC Herbicide may be mixed with pesticides as recommended by dicamba or other similar sprays, which should preferably follow FARMALINX DIMOP 375 EC Herbicide. 3. Allow at least 10 days between any application of FARMALINX DIMOP 375 EC Herbicide and other herbicides containing 2, 4-D, MCPA, Bromoxynil + MCPA at rates above 1.4 L/ha when mixing with FARMALINX DIMOP 375 EC Herbicide. 2. When mixing with broadleaf herbicides and a non-ionic surfactant (1000 g/L) at the rate of 250 mL per 100 L of spray solution. Reductions in grass weed control (as indicated above) can be minimised by using a minimum of 80 L of water per hectare by ground and 30 L by air, and if Wild Oats are present, use 2.0 L FARMALINX DIMOP 375 EC Herbicide/ha (except barley). DO NOT use

CONDITIONS OF SALE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

* Other Trademarks

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Dimop 375 EC

HERBICIDE

ACTIVE CONSTITUENT: 375 g/L DICLOFOP-METHYL
SOLVENT: 474 g/L LIQUID HYDROCARBONS

GROUP A HERBICIDE

For the post-emergent herbicide control of annual ryegrass, common barbigrass and wild oats in wheat, linseed, peas and other crops and for the control of crowsfoot grass in turf as specified in the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bond Junction NSW 2022

farmalinx

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2,
101 Grafton Street, Bondi Junction NSW 2022
T: 02 9389 2455 F: 02 9389 2844
Website: www.farmalinx.com

APVMA Approval No. 70403/63383