

OxaFert

HERBICIDE AND FERTILISER

DIRECTIONS FOR USE

RESTRAINTS:

- DO NOT mix granules with the soil.
- DO NOT apply granules when the turf foliage is wet.
- DO NOT use on putting greens, low cut tees or bowling greens.
- DO NOT apply to newly seeded turf areas.
- DO NOT apply to soft, annual flowering plants, ferns or bulbs unless they are listed as tolerant species below.
- DO NOT use in glass houses or poorly ventilated polyhouses.

Situation	Target	Turf Species	Rate	Critical Comments
Warm season turfgrass areas, including fairways, golf tees, municipal sports fields, industrial and commercial lawns	Pre-emergent control of Summer Grass (<i>Digitaria ciliaris</i>), Crowsfoot Grass (<i>Elcusine indica</i>)	Kikuyu (<i>Pennisetum clandestinum</i>), Buffalo Grass (<i>Stenotaphrum secundatum</i>), Common Couch (<i>Cynodon dactylon</i>), <i>C. transvaalensis</i>)	300 kg/ha (3 kg/100 m ²)	The product is ready to apply. No mixing or dilution is required. Apply product evenly to dry foliage and bare areas using a granular spreader able to give even coverage. Application must be made before weeds have begun to germinate. Established weeds will not be controlled. After application water in with at least 10mm of sprinkler irrigation. Control of germinating weed seedlings can be expected for up to 10 weeks after application depending upon temperature, rainfall, soil type and other factors. If longer control is required a repeat application should be made 7-10 weeks after initial application. DO NOT mow treated area until product has been watered in and turf foliage has dried. Avoid application when daytime temperatures are consistently above 32°C. This product is not recommended for use on putting greens, low cut tees or bowling greens. DO NOT apply on turf containing Carpet Grass (<i>Axonopus</i> spp.), Centipede Grass (<i>Eremochloa ophiuroides</i>) and Dichondra unless some damage is acceptable.
	Pre-emergent Control of Winter Grass (<i>Poa annua</i>) Creeping Oxalis (<i>Oxalis corniculata</i>)	Grass (Not Santa Ana) Tif Hybrid Couch Grass, Saltwater couch (<i>Paspalum vaginatum</i>), Queensland Blue Couch (<i>Digitaria didactyla</i>), Bahla grass (<i>Paspalum notatum</i>), Zoysia grass (<i>Zoysia matrella</i>)	400 kg/ha (4 kg/100 m ²)	
	Turf nutrition – provides slow release nitrogen for 8 weeks feeding		300-400 kg/ha (3-4 kg/100m ²)	

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Container-grown woody ornamental shrubs, trees and vines. (See list of tolerant species)	Bittercress (Flickweed) (<i>Cardamine hirsuta</i>) Creeping Oxalis (Yellow Wood Sorrel) (<i>Oxalis corniculata</i>) Petty Spurge (<i>Euphorbia pepulus</i>) Sorrel (<i>Rumex acetosella</i>) Sowthistle (<i>Sonchus</i> spp.) Trefloils (Lotus) (<i>Lotus</i> spp.) Willow Herbs (<i>Epilobium</i> spp.) Winter Grass (Goosegrass) (<i>Poa annua</i>)	400 kg per ha	Apply to soil surface of newly potted or freshly weeded containers. Repeat applications may be made at 10 week intervals.
		or 40 g/m ²	
Field-grown ornamental crops Transplants only: Aster, Carnation, Chrysanthemum and seed and Transplant Gypsophila (NSW only)			
			Apply to freshly weeded flower beds. Repeat applications at 10 week intervals. DO NOT apply to Zinnia or annual Helichrysum seed or transplants.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

OxaFert Herbicide and Fertiliser is a convenient ready to use formulation for the control of certain annual grass weeds in turf, container grown woody ornamental shrubs, trees and vines and field grown ornamental crops and also for fertilisation where nitrogen, potassium and sulphur are required. The product should only be used when both functions are required.

NOTE TO ORNAMENTALS:

OxaFert has been used on a wide range of ornamental plant species without damage. However some species and varieties are particularly sensitive to chemical products. As this sensitivity could be related to local conditions, it is advisable to treat only a small number of plants first, in order to determine their reaction to the product, before treating the whole crop.

Tolerant Nursery Species: Some of the species of the genera listed below have been shown to tolerate OxaFert at a dose rate of 40 g per square meter.

Abelia	Araucaria	Callitris	Cupressus	Gardenia	Hibbertia	Lantana	Melaleuca	Photinia	Schefflera
Acacia	Azalea	Camellia	Dampiera	Grevillea	Ixora	Lavandula	Metrosideros	Prostanthera	Sollya
Banksia	Banksia	Chamelaucium	Dodonaea	Hakea	Jasminium	Leea	Monstera	Protea	Viburnum
Agapanthus	Bauhinia	Coleonema	Eriostemon	Hebe	Juniperus	Leptospermum	Nerium	Ravenala	Weigela
Agonis	Boronia	Coprosmia	Escallonia	Helichrysum	Kennedia	Leucodendron	Omolanthus	Rosa	Westringia
Allamanda	Browallia	Cotoneaster	Eucalyptus	(perennial)	Kunzea	Leucospermum	Pandorea	Rosmarinus	
Ardisia	Callistemon	Cuphea	Eutaxia	Hemiandra	Lagerstroemia	Macrozamia	Phoenix	Russelia	

APPLICATION

Apply using a properly calibrated granular applicator which will apply the granules in a uniform pattern. Uniform application is essential for satisfactory weed control. Calibrate the applicator prior to use according to the manufacturer's instructions, and check applicator frequently to ensure the equipment is working properly. After application, ensure granules are washed or brushed off the foliage, particularly the foliage of prostrate and dense ground-cover plants. DO NOT apply the granules when the foliage is wet.

TURF:

For a granular spreader which can be calibrated for even output and capable of giving even application. Use a double application technique: i.e. half rate width, to achieve a full rate application. Apply under calm conditions, thereby reducing drift to non-target areas and optimizing particle distribution for good weed control. Ensure turf area is free of leaves and other debris, mowing if necessary so that the granules reach the soil surface. Delay reseeding of Ryegrass or Bentgrass into treated areas for 4-5 months (Ryegrass 4 months & Bentgrass 5 months). **DO NOT** over apply, especially under the branch spread of trees and shrubs. Avoid contact with flowers and shrubs.

Suggested Spreader Settings

The suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product.

Spreader	Ground Speed	Width of coverage	Spreader Settings	
			300 kg/ha	400 kg/ha
Scotts AccuPro 2000, SR-2000 (cone 4)	4.8 km/hr	4.0 m	0 1/2	Q
Scotts R-8, R-8A (cone 4)	4.8 km/hr	4.0 m	0	P 1/2
Spyker 288 Super	4.8 km/hr	3.0 m	3.9	4.2
Vicon (O3 Series - Standard Spout)	7.3 km/hr	11.0 m	37	44

HERBICIDE RESISTANCE WARNING

OxaFert Herbicide and Fertiliser is a member of the Oxadiazole Group of herbicide. The herbicide is an inhibitor of protoporphyrinogen oxidase. For weed resistance management, the product is a Group G Herbicide. Some naturally occurring weed biotypes resistant to the herbicide may exist through normal genetic variability in weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other oxadiazole group herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Farmalinx Pty Ltd does not accept liability for any losses that may result from failure of this product to control resistant weeds.

GROUP G HERBICIDE

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to Fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply product if rapid storm run-off from treated areas into water bodies is likely to occur within several days of application.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Do not store with seed, feed, foodstuffs, insecticides, fungicides or other fertilisers. Shake bag empty into applicator. Do not dispose of unused product on site. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SPILLS

Sweep up spilled materials and return to original container. Wash contaminated equipment taking care the washings DO NOT enter sewers, drains and other waterways.

SAFETY DIRECTIONS

Avoid contact with eyes. Do not inhale dust. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for OxaFert Herbicide and Fertiliser is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE TO BUYER

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

OxaFert

HERBICIDE AND FERTILISER

ACTIVE CONSTITUENT: 10 g/kg OXADIAZON

SOLVENT: 10.5 g/kg POLYBUTENE

GROUP G HERBICIDE

Combination herbicide and fertiliser for pre-emergent control of summer grass & crab grass and winter grass in turf, feeding of turf and for the control of certain weeds in woody ornamental shrubs in nurseries.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

farmalinx

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2,
101 Grafton Street, Bondi Junction NSW 2022
T: 02 9389 2455 F: 02 9389 2844
Website: www.farmalinx.com

APVMA Approval No.: 70376/105652