

**DIRECTIONS FOR USE****Restraints**

DO NOT spray if weeds are wet.  
 DO NOT apply if rainfall is expected before spray deposits dry on plant surfaces.  
 DO NOT water treated area for 24 hours to achieve maximum results.  
 DO NOT apply if heavy rains or storms that are likely to cause runoff from soil are forecast within 3 days of application.  
 DO NOT irrigate to the point of runoff from soil for at least 3 days after application.  
 DO NOT allow spray to contact or drift onto plants you do not want killed.  
 For professional use only.

SITUATION	WEEDS	RATE	CRITICAL COMMENTS
Paths, Driveways, around sheds Gardens, Amenity horticulture areas, Protected cropping situations Around Nursery stock. Spot Spraying in Lawns and Turf.	Seedling and young annual and perennial weeds and grasses including barnyard grass, camel melon, capeweed, chickweed, common storksbill, couch grass, clover, fat hen, giant brome grass, Indian Hedge mustard, lambs tongue, peppercress, potato weed, perennial ryegrass, sowthistle, stinking roger, whorled pigeon grass.	70 mL per litre of water	Spray to completely cover weed surface. Reapplication at around 7 days may be necessary if regrowth occurs. This product may damage lawn and turf when used as a spot spray. Perennial weed species may require repeated applications to obtain long term control. To control weeds in Protected cropping situations apply on each side of the crop row taking care not to allow spray to contact any of the crop foliage.
Orchards and vineyards Fallow soil	More advanced or established weeds may only be suppressed and require repeated application.	Mix 7 L per 100 litres water. Apply 900 - 1400 L spray per ha	Apply as a band spray 400 mm wide on each side of the orchard or vine crop row. Apply at early vegetative stage of weeds. Best results are obtained at 4- 8 leaf stage when weed canopy is less than 15 cm in height. Established weeds may require more than one application. Repeat spray after 7 days if required. Spray over all weed foliage to ensure full foliage penetration and coverage. Partial foliage penetration or coverage will result in partial control.
Forestry		7 L per 100 litres water.	To assist in site preparation apply at early vegetative stage of weeds. Best results are obtained at 4- 8 leaf stage when weed canopy is less than 15 cm in height. Established weeds may require more than one application. Repeat spray after 7 days if required. Spray over all weed foliage to ensure full foliage penetration and coverage. Partial foliage penetration or coverage will result in partial control.
Gardens, Paths, Rockeries, Tiled Roofs, Walls, Driveways, Around buildings, Spot Spraying in Lawns and Turf	Algae, Lichen, Liverwort, Moss	70 mL per litre of water	Spray to completely cover plant surface. For drives or heavy infestations, clean off by brushing after 3 days and reapply if necessary.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS  
 AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS  
 NOT REQUIRED WHEN USED AS DIRECTED.**

**GENERAL INSTRUCTIONS**

Farmalinx BRUT Herbicide is a natural contact herbicide containing pelargonic acid (nonanoic acid). Pelargonic acid occurs naturally in plants including geraniums. It acts on the foliage, causing it to quickly dehydrate and collapse. Results are seen as browning and wilting usually within hours of application. To be effective, Farmalinx BRUT Herbicide must come into contact with the green parts of the plant. Farmalinx BRUT Herbicide will control many weed species and is non selective.

**MIXING**

Half fill spray tank and commence agitation. Slowly add required amount of Farmalinx BRUT Herbicide. Top up with required quantity of water. Agitate well before commencing spray procedure. Do not mix more than is needed.

**APPLICATION**

Ensure that all weed foliage is totally covered with spray as Farmalinx BRUT Herbicide is a contact spray only. Partial coverage will give only partial control. Apply by ground equipment only using for example boom or spot sprayers. Apply using a coarse droplet spray. Optimum application volumes are related to weed maturity and density. Contact with the weeds is essential for control, as such apply in sufficient water to ensure complete and thorough coverage of foliage.  
 DO NOT spray porous surfaces without prior testing. May cause discolouration on porous hard surfaces.

**PRECAUTIONS**

Wear eye protection when mixing or using.

**Re-Entry Period**

Do not allow entry into treated areas until spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:**

DO NOT contaminate wetlands or watercourses with this product or used containers. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in noncrop areas.

**SDS**

Additional information is listed in the safety data sheet (SDS). A safety data sheet for Farmalinx BRUT Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

**CONDITIONS OF SALE:**

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

**For Refillable Containers:**

Empty contents fully into application equipment. Close all valves and return to point of supply/designated collection point for refill or storage.

**SAFETY DIRECTIONS**

Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and goggles. When using the prepared spray wear gauntlet-length chemical resistant gloves, face shield or goggles. Wash hands after use. After each day's use wash gloves, face shield or goggles, and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

*farmalinx*  
 Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2,  
 101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval NO.: 89257/124182

**CAUTION**  
 KEEP OUT OF REACH OF CHILDREN  
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Brut**  
 FARMALINX

**HERBICIDE**

ACTIVE CONSTITUENT: 525 g/L NONANOIC ACID

For non-selective control of seedling and young  
 broadleaf and grass weeds, for suppression of  
 established weeds and perennial species and for  
 control of moss and algae as per the directions for use.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE  
 CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

*farmalinx*

FARMALINX Pty Ltd ABN 95 134 353 245  
 Level 25, Suite 2506, Tower 2, 101 Grafton Street,  
 Bondi Junction NSW 2022