



DIRECTIONS FOR USE:

Refer to LIST OF PLANT SPECIES for species on which this product may be used.

SITUATION	RATE	CRITICAL COMMENTS
Seed growing	1.5 mL in 1 L water	Apply 2 L of mixed solution per square metre by suitable means e.g. watering can, immediately after sowing and covering.
Pricking-off seedlings into boxes and pots	(15 mL in 10 L water)	Apply 2 L of mixed solution per square metre by suitable means e.g. watering can and leave until soil mixture surface is dry enough to prick-off.
Treatment of cuttings		Dip cuttings in mixed solution prior to planting. Apply 2 L of mixed solution per square metre of soil immediately after setting of cuttings and at intervals of 3 weeks thereafter. A total of 3 to 4 applications during the growing season should be sufficient.
Pot plants		Dip young plants prior to potting out. Apply 100 mL of mixed solution per 100 to 110 mm pot; 150 mL per 120 to 130 mm pot and repeat at intervals of 3 weeks thereafter. A total number of 3 to 4 applications during the growing season should be sufficient.
Bulk soil treatment - preventive treatments	250 to 300 mL in 10 to 20 L water per cubic metre of soil.	Apply to bulk soil for use with pot plants.
Recreational turf	45 – 65 mL/100 m ²	Apply in sufficient water to obtain coverage. Use the lower rate as a preventative treatment. Use the higher rate as a curative or as a preventative treatment during times of high infection pressure. Repeat at 3 – 4 week intervals.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THE LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

TURF: DO NOT GRAZE TREATED AREAS OR FEED LAWN (TURF) CLIPPINGS FROM TREATED AREAS TO ANIMALS, INCLUDING POULTRY

LIST OF PLANT SPECIES:

Farmalinx AVENGER Fungicide has been successfully used at the recommended dosages on the following plants:

Adiantum (pot plants)
Aechmea spp.
Aeschynanthus spp.
 Ageratum (seed sowing)
 Alyssum
 Anemone (see Ranunculus)
Anthurium spp.
 Antirrhinum (snapdragon)
 Aphelandra dania (pot plants)
 Aralia
 Asparagus
 Aster (see Callistephus)
 Azalea
Banksia spp.
Begonia spp. (also seed sowing)
Boronia spp.
 Bromeliads
Brunfelsia calysina
 Calceolaria (also see sowing)
 Calendula
 Callistemon
 Callistephus (summer aster) (seed sowing)
 Camation
 Chlorophytum
Chrysanthemum indicum (also cuttings)
Chrysanthemum segetum (seed sowing)
 Cineraria (see Senecio)
 Clerodendron
 Clematis
Codiaeum variegatum
 Coleus

Cyclamen
 Delphinium
 Dianthus
 Dieffenbachia
Dracaena spp.
Erica gracilis
Eucalyptus spp.
Euphorbia pulcherrima spp.
Fatsyhedera lizei
Fatsia japonica
Ficus spp.
Fittonia verschaffeltii
 Fuchsia
Gerbera jamesonii
 Gloxinia
Grevillea spp.
 Gynura
Hedera helix (ivy)
 Helichrysum (immortelle)
 Impatiens
 Iris
 Kalanchoe
 Leptospermum
 Lilium
 Lobelia (seed sowing)
 Maranta
 Matthiola(stock)(seed sowing)
Melaleuca spp.
Mesembryanthemum
Mikania appifolia
Monstera deliciosa

Nemesia
 Orchids (all common species)
Oxalis hedysacharoides rubra
Pachystachys spp.
Peperomia spp.
 Petunia (seed sowing)
 Phalaenopsis
Philodendron spp.
 Phlox
Pilea spp.
 Polypodium (pot plants)
 Polyscias (lace aralia)
Primula spp.
 Ranunculus (anemone)
 Rosa (seed sowing)
 Saintpaulia
 Scindapsus or
 Rhabdiphora
Senecio cruentus
Solanum capsicastrum
 Streptocarpus
 Strobilanthes
 Syngonium
 Tagetes (seed sowing)
Tristania spp.
 Tulipa (tulip)
 Verbena
 Viola tricolour (pansy) (also seed sowing)
Vriesea splendens
 Zinnia (seed sowing)

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

Farmalinx AVENGER Fungicide is a member of the carbamates group of fungicides. For fungicide resistance management AVENGER is a Group 28 fungicide. Some naturally occurring fungal populations resistant to AVENGER and other Group 28 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by AVENGER and other Group 28 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of AVENGER to control resistant fungi.

Mixing

This product is completely soluble in water. Simply add the concentrate to the water and stir thoroughly when mixing. The mixture is stable and can be used the following day.

Application Equipment

The product can be applied using a watering can or injection equipment (e.g. Gewa) or any conventional high volume ground spraying equipment calibrated to low pressure pump output. Spray nozzles should be capable of delivering a coarse high volume spray at low pressure, 50 to 100 kPa.

Compatibility

The product is compatible with most insecticides and fungicides. It is NOT compatible with foliar nutrients.

PRECAUTION

AVENGER is corrosive to metals and spray residues should be washed out of the spray vat and the spray system with clean water immediately after use.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.



FARMALINX Pty Ltd ABN 95 134 353 245
 Level 25, Suite 2506, Tower 2,
 101 Grafton Street, Bondi Junction NSW 2022
 T: 02 9389 2455 F: 02 9389 2844
 Website: www.farmalinx.com

APVMA Approval NO.: 89397/127231

CAUTION
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT:
 600 g/L PROPAMOCARB MONOHYDROCHLORIDE

GROUP 28 FUNGICIDE

A systemic fungicide for the control of damping-off in ornamental plants, turf and eucalyptus forestry production nurseries caused by Pythium spp., as indicated in the DIRECTIONS FOR USE table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.



FARMALINX Pty Ltd ABN 95 134 353 245
 Level 25, Suite 2506, Tower 2, 101 Grafton Street,
 Bondi Junction NSW 2022