

FARMALINX

Bifentin 2G

INSECTICIDE

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply inside buildings.

DO NOT apply within 2 metres of water (eg ponds, creeks, dams, drains).

SITUATION	PEST	RATE g/10 m ²	CRITICAL COMMENTS
Turf (For example: lawns, commercial turf farms, parks, recreational areas, golf courses, bowling greens, sports fields)	Lawn Armyworm (<i>Spodoptera mauritia</i>), Sod Webworm (<i>Herpetogramma licarsialis</i>)	60	Broadcast BIFENTIN 2G with suitable application equipment to ensure uniform coverage over the treatment area. To ensure optimum control, irrigate the treated area with up to 4 mm of water immediately after application.
	Argentine stem weevil adults (<i>Listronotus bonariensis</i>)	60-120	Inspect treated areas for continuing activity. Reapply as required. Where a rate range is indicated use lower rates under lower insect pressure and higher rates under higher insect pressure.
	African black beetle (<i>Heteronychus arator</i>)	120-180	
	Billbug adults (<i>Sphenophorus brunnipennis</i>)	60-120	
External surrounds of buildings and structures	Black ants, Coastal Brown Ants, Funnel Ants, Meat Ants, Sugar Ants, Stinging Ants only	60-220	Broadcast BIFENTIN 2G with suitable application equipment to ensure uniform coverage over the treatment area. Apply granules to areas where ants are active. Where possible apply granules directly to the nest. Use the low rate for maintenance treatments or to control light infestations. Use the high rate for heavy infestations and for maximum residual control. The elimination of funnel ants from a particular site will generally require more than one application. Initial applications should be broadcast over affected areas. As the initial numbers of active colonies is reduced, application should shift to targeting active mounds. Apply granules directly to the mound and in the area immediately surrounding active mounds (300 mm radius).
	Fleas (Adults and Larvae) (<i>Ctenocephalides</i> spp.)	110-220	Apply granules to areas where ants are active. Where possible apply granules directly to the nest. Use the low rate for maintenance treatments or to control light infestations. Use the high rate for heavy infestations and for maximum residual control. Perimeter Application: Where perimeter pest control is being implemented apply BIFENTIN 2G in a two or three metre band, around and adjacent to the building or structure. Pad mounted electrical sites: To reduce or prevent ant mound building activity on electrical pads, uniformly distribute 60 to 220 grams BIFENTIN 2G per 10 m ² of mounting pad. When existing transformers are completely or partially filled with soil excavated by resident ants, remove any excess fill and level the remaining soil prior to application.
	Ticks (<i>Ixodes</i> spp.)	110-220	Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. For optimum results irrigate the treated areas with up to 10 mm of water immediately after application. Use higher application rates when treating areas with dense ground cover or heavy leaf litter. Applications should be made in spring to control larvae and nymphs that reside in the soil and leaf litter. Do not make spot applications. Treat the entire area where exposure to ticks may occur. Use higher application rates when treating areas with dense ground cover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals. Retreat as necessary to maintain adequate control.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**GENERAL INSTRUCTIONS**

Farmalinx BIFENTIN 2G Insecticide is a convenient ready to use formulation for the control of the nominated pests.

For turf application use a granular spreader that can be calibrated for even output and is capable of giving even application. Use a double application technique i.e. Half rate – half width to achieve a full rate application. Apply under calm conditions, thereby reducing drift to non-target areas and optimising particle distribution for good insect control. BIFENTIN 2G can be used as a protective treatment when applied at regular intervals or as a knockdown treatment to control existing pests. Best results are obtained when BIFENTIN 2G is applied before pest populations build up to damaging levels.

PRECAUTIONS

DO NOT spread granules directly onto humans, pets, exposed food, food preparation areas or food utensils. Keep away from domestic pets, especially dogs. If dogs eat the granules, contact your veterinary surgeon immediately. Use with caution around pets - ensure product is well dispersed ie. swept between pavers, cracks and crevices and dispersed into soil and/or lawn. Cover or remove fishponds, aquariums etc before application. DO NOT apply within 2 metres of a fishpond or water containing fish or aquatic species. DO NOT apply granules directly into drains, sewers, streams or ponds.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, dry place out of reach of children. DO NOT store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS

Wash hands after use

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 0800 764 766.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for Farmalinx BIFENTIN 2G Insecticide is available from Farmalinx Pty Ltd on request. Call Customer Service on 02 9389 2455.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Bifentin 2G

FARMALINX

INSECTICIDE

ACTIVE CONSTITUENT: 2 g/kg BIFENTHRIN

For the control of certain pests in turf; ants, fleas and ticks in external surrounds of buildings and structures as specified in the Directions for Use Table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022**farmalinx****farmalinx**
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2,

101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval NO.: 88530/121613