

**DIRECTIONS FOR USE:**

**RESTRAINTS:** DO NOT apply If rain is likely before the spray is dry.

DO NOT apply to wet foliage.

DO NOT apply this product when hot conditions (>35°C) or frosts occur as damage may result.

DO NOT apply this product to copper sensitive crops or cultivars.

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Grape vines	Downy Mildew ( <i>Plasmopara viticola</i> )	<b>Dilute spraying</b> 300 to 350 g/100 L water  <b>Concentrate spraying</b> Refer to Mixing/Application section	14 days	If Downy Mildew is expected begin spray application at 10-15 cm stage of shoot growth. Continue a programme of application every 10 to 14 days until the risk of infection diminishes. Use the higher when infection levels are particularly high. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Mangoes	Anthrachnose ( <i>Collectotrichum gloeosporioides</i> )	<b>Dilute spraying</b> 300 g/100 L water  <b>Concentrate spraying</b> Refer to Mixing/Application section	14 days	Spray at weekly intervals during flowering then at two to three week intervals until harvest. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS (WHP)**

**GRAPE VINES: DO NOT HARVEST FOR 14-DAYS AFTER APPLICATION.**

**MANGOES: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.**

**GENERAL INSTRUCTIONS**

FARMALINX Copman Fungicide is a protectant fungicide for the control of certain fungus diseases in grape vines and mangoes.

**MIXING**

Add the required quantity of FARMALINX Copman Fungicide to water in the spray tank and mix thoroughly. Maintain agitation during mixing and application.

**APPLICATION****Dilute Spraying**

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the point to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

**Concentrate Spraying**

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applied water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can be then calculated in the following way:

**EXAMPLE ONLY**

- Dilute spray volume as determined above: For example 1000 L/ha
  - Your chosen concentrate spray volume: For example 250 L/ha
  - The concentration factor in this example is:  $4 \times$  (i.e.  $1000 \text{ L}/250 \text{ L} = 4$ )
  - If the dilute label rate is 300 g/100 L, then the concentrate rate becomes  $4 \times 300$ , that is 1.2 kg/100 L of concentrate spray.
- The chosen spray volume, amount of product per 100 L of water and the sprayer set up and operation may need to be changed as the crop grows.
  - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake competency training and follow industry Best Practices.

**COMPATIBILITY**

This product is compatible with formulations of most commonly used fungicides, insecticides and miticides. Check compatibility before using by preparing a test solution of the required proportions and observing stability over several minutes.

**FUNGICIDE RESISTANCE WARNING**

FARMALINX Copman Fungicide is a **GROUP M1M3 FUNGICIDE** combination of two multi-site activity fungicides. For fungicide resistance management FARMALINX Copman Fungicide is both a Group M1 and a Group M3 fungicide. Some naturally occurring individual fungi resistant to FARMALINX Copman Fungicide may exist through normal genetic variability in any fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by FARMALINX Copman Fungicide and other Group M1 or M3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of FARMALINX Copman Fungicide to control resistant fungi.

**Re-entry:** DO NOT allow entry into treated areas for 12 hours after spraying. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions or from spraying equipment, that may cause spray drift or run-off onto sensitive areas including (but not limited to) streams, rivers or waterways. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**PROTECTION OF LIVESTOCK**

DO NOT feed treated crops and or trash to livestock

**PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT**

Dangerous to fish and other aquatic organisms. DO NOT allow spray drift or run-off onto sensitive areas including (but not limited to) streams, rivers or waterways. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

**farmalinx**  
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245  
South Sydney Corporate Park, Unit W2E,  
75-85 O'Riordan Street, Alexandria NSW 2015  
T: 02 8338 9473 F: 02 9313 4352

Website: www.farmalinx.com

APVMA Approval No.: 64403/0809

**STORAGE AND DISPOSAL**

Store in the closed, original container, in a well-ventilated area, building or room, as cool as possible, away from children, animals or fodder. Keep away from foodstuffs. DO NOT store for prolonged periods in direct sunlight. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on site. Puncture or shred or bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

**SAFETY DIRECTIONS**

Will damage eyes. Will irritate nose, throat and skin. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. If product in eyes, wash it out immediately with water. When opening the container and preparing the spray, wear elbow-length nitrile or PVC gloves, goggles and disposable dust mask. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length nitrile or PVC gloves or goggles. Wash hands after use. After each days use, wash gloves, goggles and contained clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Avoid giving alcohol. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Copman Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 8338 9473.

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

**Copman**

FUNGICIDE

ACTIVE CONSTITUENTS: 120 g/kg COPPER present as the TRIBASIC SULPHATE 300 g/kg MANCOZEB

**GROUP M1M3 FUNGICIDE**

*For the control of Downy mildew in grape vines and Anthracnose in mango as shown in the Directions for Use table.*

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

FARMALINX Pty Ltd ABN 95 134 353 245  
South Sydney Corporate Park, Unit W2E,  
75-85 O'Riordan Street, Alexandria NSW 2015

**farmalinx**