

FrontRow 500 SC

HERBICIDE

DIRECTIONS FOR USE

Crops	Weeds Controlled	State	Rate L/ha	Critical Comments	
Lettuce (Direct Sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Ryegrass, Summer Grass,	All States	4.5 in 330 - 550L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
Transplanted	Wireweed, Prince of Wales Feather, Shepherds Purse, Nettles			Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
		Vic only	2.3 in 300 - 550L	Apply on light sandy soils, low in organic matter.	
Turf & Lawns Common Couch	Winter Grass	All States	1.2 (12mL/100m ²)	Rate/ Bowling Green 37 x 37m	Apply during the Autumn and Winter, if established turf is infested with seedling Winter Grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Queensland Blue Couch after 4 weeks. DO NOT RESEED FOR 60 DAYS AFTER TREATMENT. Use ONLY on established turf. DO NOT use on Bluegrass, Ryegrass, Fescue or Bent Grass Lawns.
				170mL	
		Qld only	0.6 (6mL/100m ²)	85mL	
Queensland Blue Couch		WA only	1.2 (12mL/100m ²)	170mL	
Buffalo Grass					
Lucerne, Clover, Medics etc., grown for Forage, Hay or Seed production	Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue and Rats Tail Fescue	NSW, ACT, SA, WA, Tas only	1 - 1.5	SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July. WEEDS: Apply when weeds are at post or pre-emergent states. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.	
	1.5 - 2				
	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, Seedling Sorrel, Blackberry Nightshade, Patterson's Curse, Seedling Wireweed				

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil. Mixing prior to pouring, shake container vigorously, then add the required quantity of FrontRow 500 SC Herbicide to water in the spray vat while stirring or with agitators in motion. DO NOT mix with other pesticides.

RESISTANT WEEDS WARNING

GROUP D HERBICIDE

FARMALINX FrontRow 500 SC Herbicide is a member of the benzamide group of herbicides with inhibitors of microtubule assembly mode of action. For weed resistance management this is a Group D Herbicide. Some naturally occurring weed biotypes

resistant to FARMALINX FrontRow 500 SC Herbicide and Group D Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FARMALINX FrontRow 500 SC Herbicide or Group D Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of FARMALINX FrontRow 500 SC Herbicide to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or an FARMALINX representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or postures.

PROTECTION OF LIVESTOCK

DO NOT graze sheep, on treated areas or feed to poultry or pigs.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX FrontRow 500 SC Herbicide is available from FARMALINX on request. Call Customer Service on (02) 9389 2455.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

FrontRow 500 SC

HERBICIDE

ACTIVE CONSTITUENT: 500 g/L PROPYAZAMIDE
Solvent: 42 g/L Ethylene Glycol

GROUP **D** HERBICIDE

For selective control of certain grasses and broadleaf weeds in Lettuce, Sports Turf, Home Lawns and Legume Seed Crops and Pastures as per the DIRECTIONS FOR USE table.

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022