

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT APPLY if weeds or turf are under stress or when frosts are imminent.

DO NOT USE if rain is expected within 4 hours.

DO NOT USE clippings as mulch around other plants or for composting.

DO NOT USE on bent grass or winter grass golf and bowling greens.

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Turf	Broadleaf weeds including; White Clover (<i>Trifolium repens</i>) Plantain (<i>Plantago lanceolata</i>) Capeweed (<i>Arctotheca calendula</i>) Cat's ear (<i>Hypochoeris radicata</i>) Bin-dii (Jo-Jo, Onehunga) (<i>Soliva sessilis</i>) Cudweed (<i>Gnaphalium spp.</i>) Creeping oxalis (<i>Oxalis corniculata</i>)	5 L in 200-500 L water/ha	Apply to actively growing weeds. Transient discolouration may occur on kikuyu and carpet grass and Queensland blue couch. Varietal differences in certain buffalo grasses (eg. ST85) may produce more pronounced effects and it is recommended that small areas be tested for turf safety before large-scale application occurs. Avoid exceeding application rates through overlapping sprayed areas. Use a surfactant for difficult to wet weeds.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHOLDING PERIOD:

DO NOT GRAZE TREATED TURF OR FEED TURF CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK

GENERAL INSTRUCTIONS

For best results the turf and weeds should be actively growing at the time of treatment. Complete weed death may take up to 6 weeks. Faster results can be expected on smaller weeds. Due to residual effects of diflufenican, overseeding treated areas within 3 months of application may inhibit establishment of the turf grass.

Pourability: If cold temperature environments (less than 5°C) are encountered during product preparation for application, the product will be difficult to pour. Should this be the case, it is recommended to shake or stir the contents of the product container for about 5 minutes prior to use.

Mixing: Shake original container before use. Agitate continuously to ensure thorough mixing before and during application. Only mix sufficient chemical for each day's work. Surfactants (if used) should be added after Farmalinx GAFF Selective Herbicide and water are mixed when the spray-tank is almost full.

Mowing: Avoid mowing during the 3 to 4 days preceding or following treatment.

Cleaning Spray Equipment: THOROUGHLY CLEAN ALL APPLICATION EQUIPMENT AFTER USE.

Rinse water should be discharged onto a designated disposal area or if unavailable onto unused land away from desirable plants and water-courses.

Partial Cleaning (before spraying tolerant turf): After using Farmalinx GAFF Selective Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank with a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

Complete cleaning (before spraying susceptible plants): After using Farmalinx GAFF Selective Herbicide empty the tank completely and drain the whole system. Thoroughly wash inside the tank with a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles, then drain. Quarter fill the tank again and add an alkaline detergent at 500 mL/100 L water and circulate through the system for at least 15 minutes. Drain, remove filters and nozzles and clean separately. Rinse inside the tank thoroughly using a pressure hose and flush system with clean water.

COMPATIBILITY: No information is available that indicates physical compatibility with other products. Tank-mixing other products with Farmalinx GAFF Selective Herbicide cannot therefore be recommended. As formulations of other manufacturers' products are beyond the control of Farmalinx Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

RESISTANT WEEDS WARNING

FARMALINX GAFF Selective Herbicide is a member of the nicotinamide, phenoxy and pyridine groups of herbicides. FARMALINX GAFF has the inhibitors of carotenoid biosynthesis and disruptor of cell growth modes of action. For weed resistance management, FARMALINX GAFF is a Group F and I herbicide. Some naturally-occurring weed biotypes resistant to FARMALINX GAFF and other Group F and I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FARMALINX GAFF or other Group F and I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of FARMALINX GAFF to control resistant weeds.

PRECAUTION: Manual Handling of Turf: Hand-weeding and transplanting should not be performed before 2 weeks after spray application, unless workers wear overalls (or equivalent clothing) and chemical resistant gloves.

RE-ENTRY STATEMENT: Do not allow entry into treated areas until the spray has dried. When prior entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Spray drift from Farmalinx GAFF application may cause injury to herbaceous plants, vegetables, fruit crops, vines, ornamentals and trees.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic life. Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

(10 L and 20 L only): Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers (1000 L only): Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage. This container remains the property of Farmalinx Pty Ltd.

NOTICE TO BUYER:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2,
101 Grafton Street, Bondi Junction NSW 2022
T: 02 9389 2455 F: 02 9389 2844
Website: www.farmalinx.com

APVMA Approval No.: 70225/62779

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Gaff
FARMALINX

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 300 g/L MCPA present as the potassium salt
20 g/L CLOPYRALID present as the potassium salt
15 g/L DIFLUFENICAN

GROUP F I HERBICIDE

For the control of certain broadleaf weeds in turf as specified in the Directions for Use.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

Envirodrums (100, 110, 120, 200 L only): Mini Bulk Returnable Container: Store the original sealed Envirodrum in a cool well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Farmalinx Pty Ltd.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. If product in eyes, wash it out immediately with water. Wash hands after use. When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and goggles. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX GAFF Selective Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.