

FARMALINX

# Imazalil 750 WG

FUNGICIDE

## DIRECTIONS FOR USE:

TREE CROP	DISEASE	RATE PER 100 L WATER	CRITICAL COMMENTS
Apples and Pears	Post-harvest storage decay: <i>Penicillium expansum</i> (Blue mould)	68 g provides a concentration of 500 ppm	Apply to fruit preferably within 24 hours of harvest. Use as a bulk dip, flood or drench for a maximum of 30 seconds. After treatment allow fruit to dry before storage.
Citrus Fruit	Post-harvest storage decays: <i>Penicillium italicum</i> (Blue mould), <i>Penicillium digitatum</i> (Green mould)		Apply to fruit preferably within 24 hours of harvest. Use as a bulk dip, flood or spray before waxing.
Rockmelon	Fusarium Fruit Rot ( <i>Fusarium</i> spp.), Sour Rot ( <i>Geotrichum candidum</i> ), Alternaria Fruit Rot ( <i>Alternaria</i> spp.), Rhizopus Soft Rot ( <i>Rhizopus stolonifer</i> ), Pink Mould Rot ( <i>Trichothecium roseum</i> )		<b>DO NOT use on rockmelons intended for export (except New Zealand).</b> Dip fruit for 45 seconds within 24 hours of harvest.
Potato Tubers	STORAGE DISEASES <i>Helminthosporium solani</i> (Silver Scurf), <i>Phoma exigua</i> (Gangrene), <i>Fusarium</i> spp. (Dry rot)	20 g to a maximum of 2 L water (15 g a.i imazalil)	Apply the solution as a mist at the rate of a maximum of 2 litres per tonne over the potatoes on the conveyor line prior to storage. IMAZALIL 750 WG will give control of Fusarium semitectum and significant reduction of dry rot caused by Fusarium trichotecidodes. For good control of Fusarium solani a tank mix with thiabendazole (TBZ) is advised.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**  
**WITHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS:

### FUNGICIDE RESISTANCE WARNING:

FARMALINX IMAZALIL 750 WG Fungicide is a member of the DMI group of fungicides. For fungicide resistance management IMAZALIL 750 WG is a Group 3 fungicide. Some naturally occurring individual fungi resistant to IMAZALIL 750 WG and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

IMAZALIL 750 WG is freely soluble in water and can be used without the addition of a wetting agent in water having a pH below 7. Use buffering agent to correct water pH to a neutral pH (ie pH = 6-7). Use IMAZALIL 750 WG before waxing. DO NOT use in waxes. At the indicated rate IMAZALIL 750 WG has curative and preventative action as well as antisporeulation properties.

### MIXING:

Add the required amount of product to water and mix thoroughly. No wetting agent is required in water having a pH below 7.

### Recharging dip solution

To maintain the required volume of dip solution in the tank, add the required volume of water and product and mix thoroughly. The dip solution should be replaced when sediment accumulates.

## COMPATIBILITY

Laboratory tests have shown this product to be physically compatible with various DPA formulations. Because of diversity of field conditions, small scale tests should be carried out in the first instance.

## PROTECTION OF LIVESTOCK

DO NOT feed treated produce or by-products to food producing animals, including poultry.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate ponds, rivers, waterways or dams with the chemical or used containers.

## STORAGE AND DISPOSAL:

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

## DIP DISPOSAL

Dispose of dip in an authorized dip disposal facility.

## SAFETY DIRECTIONS

Harmful if swallowed. Avoid contact with eyes and skin. Wash hands after use.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**farmalinx**  
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2,  
101 Grafton Street, Bondi Junction NSW 2022  
T: 02 9389 2455 F: 02 9389 2844  
Website: www.farmalinx.com

APVMA Approval NO.: 65935/52355

## SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for IMAZALIL 750 WG FUNGICIDE is available from FARMALINX on request. Call Customer Service on (02) 9389 2455.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

# Imazalil 750 WG

FUNGICIDE

ACTIVE CONSTITUENTS: 750 g/kg IMAZALIL present as the SULPHATE

GROUP 3 FUNGICIDE

For control of post-harvest disease of citrus, apples, pears and storage diseases in potatoes and rockmelons as specified in the Directions for Use table.

IMPORTANT: READ THIS BOOKLET BEFORE USING THIS PRODUCT

**farmalinx**

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2, 101 Grafton Street,  
Bondi Junction NSW 2022