

FARMALINX

# Propyzamide 500 WG

HERBICIDE

## DIRECTIONS FOR USE

Crops	Weeds Controlled	State	Rate kg/ha	Critical Comments
Lettuce (Direct Sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Ryegrass, Summer Grass,	All States	4.5 in 330 - 550L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.
Transplanted	Wireweed, Prince of Wales Feather, Shepherds Purse, Nettles			Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.
		Vic only	2.3 in 300 - 550L	Apply on light sandy soils, low in organic matter.
Turf & Lawns Common Couch	Winter Grass	All States	1.2 (12g/100m <sup>2</sup> )	Rate/ Bowling Green 37 x 37m 170g Apply during the Autumn and Winter, if established turf is infested with seedling Winter Grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Queensland Blue Couch after 4 weeks. <b>DO NOT RESEED FOR 60 DAYS AFTER TREATMENT.</b> Use <b>ONLY</b> on established turf. <b>DO NOT</b> use on Bluegrass, Ryegrass, Fescue or Bent Grass Lawns.
Queensland Blue Couch		Qld only	0.6 (6g/100m <sup>2</sup> )	85g
Buffalo Grass		WA only	1.2 (12g/100m <sup>2</sup> )	170g
Lucerne, Clover, Medics etc., grown for Forage, Hay or Seed production	Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue and Rats Tail Fescue	NSW, ACT, SA, WA, Tas only	1 - 1.5	<b>SEEDLING CROPS:</b> Apply after the third trifoliate leaf is fully developed. <b>ESTABLISHED CROPS:</b> Apply any time before the end of July. <b>WEEDS:</b> Apply when weeds are at post or pre-emergent states.  <b>BEST RESULTS:</b> When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting <b>PRIOR</b> to treatment is recommended.
	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, Seedling Sorrel, Blackberry Nightshade, Patterson's Curse, Seedling Wireweed		1.5 - 2	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.**

### GENERAL INSTRUCTIONS

Best results are obtained by applying Propyzamide 500 WG Herbicide to cool, wet soil.

### RESISTANT WEEDS WARNING

FARMALINX Propyzamide 500 WG Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is a Group **D** Herbicide. Some naturally occurring weed biotypes resistant to FARMALINX Propyzamide 500 WG Herbicide and Group **D** Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FARMALINX Propyzamide 500 WG Herbicide or Group **D** Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of FARMALINX Propyzamide 500 WG Herbicide to control the resistant weeds.

GROUP **D** HERBICIDE

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or an FARMALINX Pty Ltd representative.

### SPRAY PREPARATION

FARMALINX Propyzamide 500 WG Herbicide is a Wettable Granule formulation to be blended with water and applied as a spray. Granule formulations are designed to be added to large volumes of water and dispersed slowly under agitation. When using larger boomsprays and unable to add granules directly to a large volume in the tank via the top of the boomspray, pre-mixing may be necessary as follows:

- Partly (30%) fill the boomspray tank with water.
  - Slowly add required amount of FARMALINX Propyzamide 500 WG Herbicide for the area to be sprayed into Granni Pot<sup>®</sup> or other pre-mixing vessel while agitating well.
  - Ensure high water volumes and **DO NOT** add to the spray tank until the pre-mix is fully dissolved. This may take several mixes to ensure the correct amount is added to the boomspray tank.
  - Add the pre-mixed FARMALINX Propyzamide 500 WG Herbicide into the boomspray tank with the agitation system operating.
  - Fill up to the correct volume with water.
  - The mixture must be kept in suspension at all times by continuous agitation.
  - Tank-mixtures should be tested for physical compatibility and efficacy before use. **DO NOT** mix with MCPA. If tank mixing, FARMALINX Propyzamide 500 WG Herbicide must be in suspension before adding other herbicides, insecticides or surfactants.
- Wash sprayer thoroughly after use.  
**DO NOT** handle the water-soluble packaging with wet hands or wet gloves. Keep dry.

### EQUIPMENT

Always ensure the sprayer is clean, using ChemAg Spray and Boom Cleaner as directed. It is recommended that for certain turf situations up to 100L/ha spray volume is required.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

**DO NOT** apply to, or allow spray to drift onto crops, pastures or non-target desirable plants, nor onto agricultural land scheduled for cropping or pasture, as injury to the crop or pasture may occur.

### PROTECTION OF LIVESTOCK

**DO NOT** graze sheep, cut for stock feed (including poultry and pigs) any treated areas (for 25 days after application).

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

**DO NOT** contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not

available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

### SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

### SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for FARMALINX Propyzamide 500 WG Herbicide is available from FARMALINX on request. Call Customer Service on (02) 9389 2455.

### Suspected of causing cancer.

If exposed or concerned: Get medical advice.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Propyzamide 500 WG**

HERBICIDE

ACTIVE CONSTITUENT: 500 g/kg PROPYZAMIDE

GROUP **D** HERBICIDE

For selective control of certain grasses and broadleaf weeds in Lettuce, Sports Turf, Home Lawns and Legume Seed Crops and Pastures as per the DIRECTIONS FOR USE table.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

**farmalinx**  
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2,  
101 Grafton Street, Bondi Junction NSW 2022  
T: 02 9389 2455 F: 02 9389 2844  
Website: www.farmalinx.com

APVMA Approval No. 65925/52306

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2, 101 Grafton Street,  
Bondi Junction NSW 2022

**farmalinx**