FARMALINX

Snooker SL

PLANT GROWTH REGULANT

DIRECTIONS FOR USE

RESTRAINTS

DO NOT use if rain is likely within 2 hours of application. DO NOT apply from aircraft if wind speed exceeds 16 km/hour.

CROP	RATE PER HECTARE	STATES	CRITICAL COMMENTS
Wheat	500 mL to 1.3 L	NSW, ACT, Vic, Tas, SA, WA only	Apply at Zadoks stage Z 25 to Z 31, using a minimum of 30 litres total spray per hectare for aerial application and 100 litres per hectare by ground rig. Use the higher rates in irrigated or potentially high-yielding crops. Addition of 1 mL non-ionic surfactant per litre of spray is desirable.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED. GRAZING: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

MIXING

Put half the required amount of hard or soft water in the spray tank, start agitation, add the recommended quantity of SNOOKER SL then add the rest of the water. Add non-ionic surfactant last.

Use of SNOOKER SL may result in improved yields of wheat. Treated plants generally have shorter, thicker stems and greater root development than untreated plants. Photosynthesis, translocation, nitrogen uptake and moisture utilisation are believed to be improved following use of SNOOKER SL. Lodging may be reduced.

SNOOKER SL Plant Growth Regulant should be applied to vigorously growing crops that have an adequate supply of soil moisture and nutrients in irrigated wheat, high levels of nitrogen supplementation together with adequate irrigation improve results with SNOOKER SL.

COMPATIBILITY

SNOOKER SL Plant Growth Regulant may be applied as spray tank mixtures with commonly used herbicides for which timing coincides eg., diclofop methyl, 2,4-D amine, MCPA, bromoxynil, dicamba, terbutryne and linuron.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spray equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers, drains or streams with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

For refillable containers, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX SNOOKER SL Plant Growth Regulant is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

POISON

NOTICE TO BUYER:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.



 FARMALINX Pty Ltd ABN 95 134 353 245

 Level 25, Suite 2506, Tower 2,

 101 Grafton Street, Bondi Junction NSW 2022

 T: 02 9389 2455

 F: 02 9389 2455

 F: 02 9389 case

 Website: www.farmalinx.com

APVMA Approval No.: 69925/62070



ACTIVE CONSTITUENT: 582 g/L CHLORMEQUAT present as the CHLORMEQUAT CHLORIDE ACALORMEQUAT CHLORIDE As an aid in increasing grain yields and

prevention of lodging in wheat. MPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER: READ THOROUGHLY BEFORE OPENING OR USING THIS PRO FARMALINX Pty Ltd ABN 95 134 353 245 Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022

