

Wildman 240 EC

HERBICIDE

DIRECTIONS FOR USE

Restrains: DO NOT apply if rainfall is expected within 2 hours.
DO NOT apply to weeds or crops which are under stress. For example, due to very dry, very wet, frosty or diseased conditions, nutrient deficiency or high insect pressure.
DO NOT apply to weeds or crops immediately after a prolonged period of frosts.
DO NOT apply under poor growing conditions.

CROP	WEEDS CONTROLLED	STATE	RATE PER HA	WEED GROWTH STAGE	CRITICAL COMMENTS
Wheat 2 leaf to late jointing (Zadocks 12 – 37)	Wild Oats	Qld, Nthn NSW only	65 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 to 5 leaf	NOT TO BE TANK MIXED WITH BROADLEAF HERBICIDES AT THESE RATES. Apply at 85 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 leaf to mid tillering	
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 to 5 leaf	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing FARMALINX Wildman 240 EC Herbicide with certain broadleaf weed herbicides some reduction in grass weed control may occur. Unless otherwise specified always use D-C-Trate, Uptake or Hasten when tank mixing broadleaf herbicides.
			125 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 leaf to mid tillering	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing FARMALINX Wildman 240 EC Herbicide with certain broadleaf weed herbicides some reduction in grass weed control may occur. Refer to compatibility chart for specific rates for tank mixes. Unless otherwise specified always use D-C-Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 85mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
		Sthn NSW, Vic, SA, WA only	65 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water.	2 to 5 leaf	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing FARMALINX Wildman 240 EC Herbicide with certain broadleaf weed herbicides some reduction in grass weed control may occur. Refer to compatibility chart for specific rates for tank mixes. Unless otherwise specified always use D-C-Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 85mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water.	2 leaf to mid tillering	

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE	WEED GROWTH STAGE	CRITICAL COMMENTS
Wheat 2 leaf to late jointing (Zadocks 12 – 37) – continued	Paradoxa Grass (<i>Phalaris paradoxa</i>), Canary Grass (<i>Phalaris minor</i>)	Qld, NSW, Vic, SA only	85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 to 3 leaf	NOT TO BE TANK MIXED WITH BROADLEAF HERBICIDES AT RATES BELOW 160 mL/ha. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing FARMALINX Wildman 240 EC Herbicide with certain broadleaf weed herbicides some reduction in grass weed control may occur. Refer to compatibility chart for specific rates for tank mixes. Unless otherwise specified always use D-C-Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 85mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
			125 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 to 5 leaf	
			160 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 leaf to early tillering	
	Annual Ryegrass	NSW, Vic, SA, WA only	160 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 to 4 leaf	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing FARMALINX Wildman 240 EC Herbicide with certain broadleaf weed herbicides some reduction in grass weed control may occur. Apply at 210mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
			210 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 leaf to early tillering	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

HARVESTED GRAIN: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.
ANIMAL GRAZING/FEEDING: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

www.farmalinx.com

CONTINUED OVERLEAF

least 15 minutes and see a doctor.
a glass of water. If in eyes, hold eyes open, flood with water for at
Phone Australia 131 126. If swallowed, do NOT induce vomiting. Give
Poisoning occurs contact a doctor or Poisons Information Centre.
FIRST AID

DO NOT apply to irrigation drains or channels. DO NOT contaminate
dams, streams, creeks and rivers with the chemical, spray mixture,
used containers or equipment washings.
ENVIRONMENT
DO NOT apply to irrigation drains or channels. DO NOT contaminate
dams, streams, creeks and rivers with the chemical, spray mixture,
used containers or equipment washings.
PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND
PASTURES. DO NOT allow spray to drift onto adjacent fallow land.
spraying equipment which could be expected to cause spray to drift
onto nearby susceptible plants, adjacent crops, crop lands or
pastures. DO NOT allow spray to drift onto adjacent fallow land.
BROADLEAF HERBICIDES: Some broadleaf weed herbicides are
incompatible with FARMALINX Wildman 240 EC Herbicide because
they reduce the effectiveness of FARMALINX Wildman 240 EC
Herbicide or because in mixture they may cause crop yellowing. For
the latest information on compatibility of FARMALINX Wildman 240 EC
Herbicide with broadleaf herbicides refer to your reseller or
FARMALINX Pty Ltd.

When FARMALINX Wildman 240 EC Herbicide is applied with
compatible herbicides always use a recommended oil. Only use a
100% non-ionic surfactant when mixing with diflufenican based
products.
Grass Herbicides: FARMALINX Wildman 240 EC Herbicide is
compatible with diclofop methyl.
BROADLEAF HERBICIDES: Some broadleaf weed herbicides are
incompatible with FARMALINX Wildman 240 EC Herbicide because
they reduce the effectiveness of FARMALINX Wildman 240 EC
Herbicide or because in mixture they may cause crop yellowing. For
the latest information on compatibility of FARMALINX Wildman 240 EC
Herbicide with broadleaf herbicides refer to your reseller or
FARMALINX Pty Ltd.

When FARMALINX Wildman 240 EC Herbicide is applied with
compatible herbicides always use a recommended oil. Only use a
100% non-ionic surfactant when mixing with diflufenican based
products.
Grass Herbicides: FARMALINX Wildman 240 EC Herbicide is
compatible with diclofop methyl.
BROADLEAF HERBICIDES: Some broadleaf weed herbicides are
incompatible with FARMALINX Wildman 240 EC Herbicide because
they reduce the effectiveness of FARMALINX Wildman 240 EC
Herbicide or because in mixture they may cause crop yellowing. For
the latest information on compatibility of FARMALINX Wildman 240 EC
Herbicide with broadleaf herbicides refer to your reseller or
FARMALINX Pty Ltd.

When FARMALINX Wildman 240 EC Herbicide is applied with
compatible herbicides always use a recommended oil. Only use a
100% non-ionic surfactant when mixing with diflufenican based
products.
Grass Herbicides: FARMALINX Wildman 240 EC Herbicide is
compatible with diclofop methyl.
BROADLEAF HERBICIDES: Some broadleaf weed herbicides are
incompatible with FARMALINX Wildman 240 EC Herbicide because
they reduce the effectiveness of FARMALINX Wildman 240 EC
Herbicide or because in mixture they may cause crop yellowing. For
the latest information on compatibility of FARMALINX Wildman 240 EC
Herbicide with broadleaf herbicides refer to your reseller or
FARMALINX Pty Ltd.

* Registered Trademarks

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2,
101 Grafton Street, Bondi Junction NSW 2022
T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval No.: 65388/52910

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Wildman 240 EC

HERBICIDE

ACTIVE CONSTITUENTS: 240 g/L CLODINAFOF-PROPRARGYL
60 g/L CLOQUINTOCET-IMEXYL
SOLVENTS: 575 g/L LIQUID HYDROCARBONS
100 g/L N-METHYL-2-PYRROLIDONE

GROUP A HERBICIDE

For the control of Wild Oats, Paradoxa Grass (Annual Phalaris), Canary Grass
and Annual Ryegrass in Wheat. Highly active on Wild Oats and can be mixed
with many broadleaf herbicides as per the Directions for Use Table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE
CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

farmalinx

GENERAL INSTRUCTIONS

FARMALINX Wildman 240 EC Herbicide should be applied only when
weeds are actively growing.

RESISTANT WEEDS WARNING

FARMALINX Wildman 240 EC Herbicide is a member of the **GROUP A HERBICIDE**
aryloxyphenoxypropionate (“fop”) group of herbicides. FARMALINX
Wildman 240 EC Herbicide has the inhibition of fat (lipid) synthesis,
(or inhibitors of acetyl CoA carboxylase) mode of action. For weed
resistance management FARMALINX Wildman 240 EC Herbicide is a
Group A herbicide. Some naturally-occurring weed biotypes resistant
to FARMALINX Wildman 240 EC Herbicide and other Group A
herbicides may exist through normal genetic variability in any weed
population. The resistant individuals can eventually dominate the
weed population if these herbicides are used repeatedly. These
resistant weeds will not be controlled by FARMALINX Wildman 240
EC Herbicide or other Group A herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to
use, FARMALINX Pty Ltd accepts no liability for any losses that may
result from the failure of FARMALINX Wildman 240 EC Herbicide to
control resistant weeds.

Weed biotypes resistant to diclofop-methyl and other “grass specific” herbicides can also often be resistant to FARMALINX Wildman 240 EC Herbicide. Before using FARMALINX Wildman 240 EC Herbicide on a population resistant to or suspected to be resistant to “grass specific” herbicides, have a resistance test conducted to ensure that it is susceptible to FARMALINX Wildman 240 EC Herbicide.

Large numbers of healthy surviving weeds can be an indication that
resistance is developing. Efforts should be made to prevent seed set
of the surviving weeds.

It is important to ensure that each individual component of the tank
mix is fully dissolved and in solution before the next product is added
to the tank mix, otherwise mixing problems may occur.
MINIMUM OIL VOLUMES
The minimum amount of oil to be applied with FARMALINX Wildman
240 EC Herbicide will depend on the type of oil used. When applying
FARMALINX Wildman 240 EC Herbicide using low water volumes, DO
NOT use less than the minimum oil volumes specified below:

- For sites normally applied at 1% v/v (eg D-C-Trate), DO NOT apply
less than 500 mL oil/ha.
- For sites normally applied at 0.5% v/v (eg Uptake, Hasten), DO NOT
apply less than 250 mL oil/ha.

CLEANING SPRAY EQUIPMENT
After using FARMALINX Wildman 240 EC Herbicide, empty the tank
completely and drain the whole system. Thoroughly wash inside the
tank using a pressure hose, drain the tank and clean any filters in the
tank, pump, line and nozzles.
To rinse: After cleaning the tank as above, quarter fill the tank with
clean water and circulate through the pump, lines, hoses and
nozzles. Drain and repeat the rinsing procedure twice.
To decontaminate: Before spraying cereals (except wheat), maize,
sorghum or other sensitive crops, wash the tank and rinse the
system as above. Drain and repeat the rinsing procedure. Then
quarter fill the tank and add a liquid alkali detergent (eg SURF OMO,
DRIVE) at 0.5 L/100 L of water and circulate throughout the system
for at least fifteen minutes. Drain the whole system. Remove filters
and nozzles and clean them separately. Finally flush the system with
clean water and allow to drain. Dispose of all water used for cleaning
in a disposal pit specifically marked and set up for this purpose clear
of waterways, vegetation and roots.

DO NOT make more than one application of a Group A herbicide with
the inhibition of fat (lipid) synthesis (or inhibitors of acetyl CoA
carboxylase) mode-of-action to a crop in the same season.
If the user suspects that the target weed population is resistant to
herbicides with this mode of action, FARMALINX Wildman 240 EC
Herbicide or other Group A herbicides should not be used.
Strategies to minimize the risk of herbicide resistance are available.
The above recommendations should be incorporated into an
Integrated Weed Management (IWM) Program. Consult your farm
chemical supplier, consultant, local Department of Agriculture or
Primary Industries, or local FARMALINX representative for details.

MIXING

Fill the spray tank to one quarter full. Then add FARMALINX Wildman
240 EC Herbicide while adding water to make up the final spray
volume. Pour FARMALINX Wildman 240 EC Herbicide into the stream
of incoming water. Good agitation at the time FARMALINX Wildman
240 EC Herbicide is added to the tank will ensure good mixing. Add
the required amount of D-C-Trate*, Hasten*, or Uptake* oil just before
the tank is full of water and with the agitators in motion.

MIXING ORDER

- Some products may react with other products if they are not mixed
in the correct order. The general mixing order of products should be:-
- water conditioners or buffers
 - water dispersible granules (WG)
 - wettable powders (WP)
 - flowable or suspension concentrates (SC)
 - emulsifiable concentrates (EC) (FARMALINX Wildman 240 EC Herbicide)
 - water based or soluble concentrates
 - oils and wetters – eg Hasten, BS 1000