

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50 m of crop or within 50 m of native vegetation. A 50m buffer zone in crops must be employed to reduce risk to birds and other non-target animals.

DO NOT apply bait unless monitoring of mouse numbers indicate the potential for crop damage of economic significance.

Situation	Pest	Rate	Critical Comments
Grain crops, legume crops, canola, safflower, nut crops and pasture. (Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seed has been sown into crop stubble or other similar ground cover).	Mice	1 kg/ha to achieve an even coverage of 1 pellet/m ²	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 1 pellet/m ² . It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals. Farmalinx ZINCPHOS Pellets will provide a high level of control in most situations. However in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying the bait and that post-baiting estimates are conducted to assess effectiveness of the control operation. For information on the most effective methods of monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

DO NOT harvest any crops or allow livestock to graze baited area for 14 days after application.

PROTECTION OF WILDLIFE THE ENVIRONMENT AND LIVESTOCK

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition.

Dispose of used containers by crushing and burying below one metre in a disposal pit specifically marked for this purpose, clear of waterways, desirable vegetation and tree roots. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using the baits wear elbow-length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use wash gloves and respirator and if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. DO NOT give direct mouth to mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one way mask. Resuscitate in a well-ventilated area.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX ZincPhos Pellets is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2,

101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval No.: 80058/100141

DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

ZincPhos

PELLETS

ACTIVE CONSTITUENT: 25 g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENCED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

farmalinx